

CHAPTER
16

ACID-BASE
EQUILIBRIA



VARIETIES OF CITRUS FRUITS
(oranges, limes, lemons,
grapefruit, and tangerines).

- 16.1 Acids and Bases: A Brief Review**
We start by reviewing the *Arrhenius* definitions of acid and base.
- 16.2 Brønsted–Lowry Acids and Bases**
We then learn the more general *Brønsted–Lowry* definitions for acid and base. A Brønsted–Lowry acid is a *proton donor* and a Brønsted–Lowry base is a *proton acceptor*. Two species that differ by the presence or absence of a proton are known as a *conjugate acid–base pair*.
- 16.3 The Autoionization of Water**
We will see that the *autoionization* of water produces small concentrations of H_3O^+ and OH^- ions. The *equilibrium constant* for autoionization, $K_w = [\text{H}_3\text{O}^+][\text{OH}^-]$, defines the relationship between H_3O^+ and OH^- concentrations in aqueous solutions.
- 16.4 The pH Scale**
We will use the pH scale ($\text{pH} = -\log[\text{H}^+]$) to describe the acidity or basicity of an aqueous solution. Neutral solutions have a $\text{pH} = 7$, acidic solutions have pH below 7, and basic solutions have pH above 7.
- 16.5 Strong Acids and Bases**
We categorize acids and bases as being either strong or weak electrolytes. *Strong* acids and bases are strong electrolytes, ionizing or dissociating completely in aqueous solution. *Weak* acids and bases are weak electrolytes and therefore ionize only partially.
- 16.6 Weak Acids**
We learn that the ionization of a weak acid in water is an equilibrium process with an equilibrium constant K_a that can be used to calculate the pH of a weak acid solution.
- 16.7 Weak Bases**
We learn that the ionization of a weak base in water is an equilibrium process with equilibrium constant K_b that can be used to calculate the pH of a weak base solution.
- 16.8 Relationship between K_a and K_b**
We will see that a relationship exists between the K_a and K_b of any conjugate *acid–base* pair: $K_a \times K_b = K_w$. Thus, the stronger an acid, the weaker its conjugate base.
- 16.9 Acid–Base Properties of Salt Solutions**
We will explore the fact that the ions of a soluble ionic compound (a salt) can serve as Brønsted–Lowry acids or bases.
- 16.10 Acid–Base Behavior and Chemical Structure**
We continue by exploring the relationship between chemical structure and acid–base behavior.
- 16.11 Lewis Acids and Bases**
Finally, we learn the *Lewis* definitions of acid and base. A Lewis acid is an *electron-pair acceptor*, and a Lewis base is an *electron-pair donor*.

WHAT IS THE SOUREST FOOD YOU'VE EVER TASTED? Citrus fruit, such as the lemons shown in the chapter-opening photograph? Sour cherries? Rhubarb? The sour taste of foods is due primarily to the presence of acids. Citric acid ($\text{H}_3\text{C}_6\text{H}_5\text{O}_7$), malic acid ($\text{H}_2\text{C}_4\text{H}_4\text{O}_5$), oxalic acid ($\text{H}_2\text{C}_2\text{O}_4$), and ascorbic

acid, also known as vitamin C ($\text{HC}_6\text{H}_7\text{O}_6$), are present in many fruits as well as in certain vegetables, such as rhubarb and tomatoes.

Acids and bases are important in numerous chemical processes that occur around us—from industrial processes to biological ones, from reactions in the laboratory to those in our environment. The time required for a metal object immersed in water to corrode, the ability of an aquatic environment to support fish and plant life, the fate of pollutants washed out of the air by rain, and even the rates of reactions that maintain our lives all critically depend upon the acidity or basicity of solutions. Indeed, an enormous amount of chemistry can be understood in terms of acid–base reactions.

We have encountered acids and bases many times in earlier discussions. For example, a portion of Chapter 4 focused on their reactions. But what makes a substance behave as an acid or as a base? In this chapter we reexamine acids and bases, taking a closer look at how they are identified and characterized. In doing so, we will consider their behavior both in terms of their structure and bonding and in terms of the chemical equilibria in which they participate.

16.1 ACIDS AND BASES: A BRIEF REVIEW


From the earliest days of experimental chemistry, scientists have recognized acids and bases by their characteristic properties. Acids have a sour taste and cause certain dyes to change color (for example, litmus turns red on contact with acids). Indeed, the word *acid* comes from the Latin word *acidus*, meaning sour or tart. Bases, in contrast, have a bitter taste and feel slippery (soap is a good example). The word *base* comes from an old English meaning of the word, which is “to bring low.” (We still use the word *debase* in this sense, meaning to lower the value of something.) When bases are added to acids, they lower the amount of acid. Indeed, when acids and bases are mixed in certain proportions, their characteristic properties disappear altogether. ∞ (Section 4.3)

Historically, chemists have sought to relate the properties of acids and bases to their compositions and molecular structures. By 1830 it was evident that all acids contain hydrogen but not all hydrogen-containing substances are acids. During the 1880s, the Swedish chemist Svante Arrhenius (1859–1927) linked acid behavior with the presence of H^+ ions and base behavior with the presence of OH^- ions in aqueous solution.

Arrhenius defined acids as substances that produce H^+ ions in water and bases as substances that produce OH^- ions in water. Indeed, the properties of aqueous solutions of acids, such as sour taste, are due to $H^+(aq)$, whereas the properties of aqueous solutions of bases are due to $OH^-(aq)$. Over time the Arrhenius concept of acids and bases came to be stated in the following way:

- An *acid* is a substance that, when dissolved in water, increases the concentration of H^+ ions.
- A *base* is a substance that, when dissolved in water, increases the concentration of OH^- ions.

Hydrogen chloride is an Arrhenius acid. Hydrogen chloride gas is highly soluble in water because of its chemical reaction with water, which produces hydrated H^+ and Cl^- ions:



The aqueous solution of HCl is known as hydrochloric acid. Concentrated hydrochloric acid is about 37% HCl by mass and is 12 M in HCl. Sodium hydroxide, on the other hand, is an Arrhenius base. Because NaOH is an ionic compound, it dissociates into Na^+ and OH^- ions when it dissolves in water, thereby releasing OH^- ions into the solution.



GIVE IT SOME THOUGHT


What two ions are central to the Arrhenius definitions of acids and bases?

16.2 BRØNSTED–LOWRY ACIDS AND BASES

The Arrhenius concept of acids and bases, while useful, has limitations. For one thing, it is restricted to aqueous solutions. In 1923 the Danish chemist Johannes Brønsted (1879–1947) and the English chemist Thomas Lowry (1874–1936) independently proposed a more general definition of acids and bases. Their concept is based on the fact that *acid–base reactions involve the transfer of H^+ ions from one substance to another.*

The H⁺ Ion in Water

In Equation 16.1 hydrogen chloride is shown ionizing in water to form H⁺(aq). An H⁺ ion is simply a proton with no surrounding valence electron. This small, positively charged particle interacts strongly with the nonbonding electron pairs of water molecules to form hydrated hydrogen ions. For example, the interaction of a proton with one water molecule forms the **hydronium ion**, H₃O⁺(aq):




The formation of hydronium ions is one of the complex features of the interaction of the H⁺ ion with liquid water. In fact, the H₃O⁺ ion can form hydrogen bonds to additional H₂O molecules to generate larger clusters of hydrated hydrogen ions, such as H₅O₂⁺ and H₉O₄⁺ (Figure 16.1 ▶).

Chemists use H⁺(aq) and H₃O⁺(aq) interchangeably to represent the same thing—namely the hydrated proton that is responsible for the characteristic properties of aqueous solutions of acids. We often use the H⁺(aq) ion for simplicity and convenience, as we did in Equation 16.1. The H₃O⁺(aq) ion, however, more closely represents reality.

Proton-Transfer Reactions

When we closely examine the reaction that occurs when HCl dissolves in water, we find that the HCl molecule actually transfers an H⁺ ion (a proton) to a water molecule as depicted in Figure 16.2 ▶. Thus, we can represent the reaction as occurring between an HCl molecule and a water molecule to form hydronium and chloride ions:




The polar H₂O molecule promotes the ionization of acids in water solution by accepting a proton to form H₃O⁺.

Brønsted and Lowry proposed definitions of acids and bases in terms of their ability to transfer protons:


- An *acid* is a substance (molecule or ion) that donates a proton to another substance.
- A *base* is a substance that accepts a proton.

Thus, when HCl dissolves in water (Equation 16.3), HCl acts as a **Brønsted–Lowry acid** (it donates a proton to H₂O), and H₂O acts as a **Brønsted–Lowry base** (it accepts a proton from HCl).


Because the emphasis in the Brønsted–Lowry concept is on proton transfer, the concept also applies to reactions that do not occur in aqueous solution. In the reaction between HCl and NH₃, for example, a proton is transferred from the acid HCl to the base NH₃:



This reaction can occur in the gas phase. The hazy film that forms on the windows of general chemistry laboratories and on glassware in the lab is largely solid NH₄Cl formed by the gas-phase reaction of HCl and NH₃ (Figure 16.3 ▼).



▲ **Figure 16.1 Hydrated hydronium ions.** Lewis structures and molecular models for H₅O₂⁺ and H₉O₄⁺. There is good experimental evidence for the existence of both these species.




▲ **Figure 16.2 A proton-transfer reaction.** When a proton is transferred from HCl to H₂O, HCl acts as the Brønsted–Lowry acid and H₂O acts as the Brønsted–Lowry base.



▲ **Figure 16.3** A gas-phase acid–base reaction. The $\text{HCl}(g)$ escaping from concentrated hydrochloric acid and the $\text{NH}_3(g)$ escaping from aqueous ammonia (here labeled ammonium hydroxide) combine to form a white fog of $\text{NH}_4\text{Cl}(s)$.

Let's consider another example that compares the relationship between the Arrhenius definitions and the Brønsted–Lowry definitions of acids and bases—an aqueous solution of ammonia, in which the following equilibrium occurs:



Ammonia is an Arrhenius base because adding it to water leads to an increase in the concentration of $\text{OH}^-(aq)$. It is a Brønsted–Lowry base because it accepts a proton from H_2O . The H_2O molecule in Equation 16.5 acts as a Brønsted–Lowry acid because it donates a proton to the NH_3 molecule.

An acid and a base always work together to transfer a proton. In other words, a substance can function as an acid only if another substance simultaneously behaves as a base. To be a Brønsted–Lowry acid, a molecule or ion must have a hydrogen atom that it can lose as an H^+ ion. To be a Brønsted–Lowry base, a molecule or ion must have a nonbonding pair of electrons that it can use to bind the H^+ ion.


Some substances can act as an acid in one reaction and as a base in another. For example, H_2O is a Brønsted–Lowry base in its reaction with HCl (Equation 16.3) and a Brønsted–Lowry acid in its reaction with NH_3 (Equation 16.5). A substance that is capable of acting as either an acid or a base is called **amphiprotic**. An amphiprotic substance acts as a base when combined with something more strongly acidic than itself and as an acid when combined with something more strongly basic than itself.

GIVE IT SOME THOUGHT

In the forward reaction, which substance acts as the Brønsted–Lowry base:
 $\text{HSO}_4^-(aq) + \text{NH}_3(aq) \rightleftharpoons \text{SO}_4^{2-}(aq) + \text{NH}_4^+(aq)$?

Conjugate Acid–Base Pairs


In any acid–base equilibrium both the forward reaction (to the right) and the reverse reaction (to the left) involve proton transfers. For example, consider the reaction of an acid, which we will denote HX , with water:



In the forward reaction HX donates a proton to H_2O . Therefore, HX is the Brønsted–Lowry acid, and H_2O is the Brønsted–Lowry base. In the reverse reaction the H_3O^+ ion donates a proton to the X^- ion, so H_3O^+ is the acid and X^- is the base. When the acid HX donates a proton, it leaves behind a substance, X^- , which can act as a base. Likewise, when H_2O acts as a base, it generates H_3O^+ , which can act as an acid.


An acid and a base such as HX and X^- that differ only in the presence or absence of a proton are called a **conjugate acid–base pair**.^{*} Every acid has a **conjugate base**, formed by removing a proton from the acid. For example, OH^- is the conjugate base of H_2O , and X^- is the conjugate base of HX . Similarly, every base has associated with it a **conjugate acid**, formed by adding a proton to the base. Thus, H_3O^+ is the conjugate acid of H_2O , and HX is the conjugate acid of X^- .

In any acid–base (proton-transfer) reaction we can identify two sets of conjugate acid–base pairs. For example, consider the reaction between nitrous acid (HNO_2) and water:



^{*}The word conjugate means “joined together as a pair.”

Likewise, for the reaction between NH_3 and H_2O (Equation 16.5), we have



SAMPLE EXERCISE 16.1 | Identifying Conjugate Acids and Bases

(a) What is the conjugate base of each of the following acids: HClO_4 , H_2S , PH_4^+ , HCO_3^- ? (b) What is the conjugate acid of each of the following bases: CN^- , SO_4^{2-} , H_2O , HCO_3^- ?

SOLUTION

Analyze: We are asked to give the conjugate base for each of a series of species and to give the conjugate acid for each of another series of species.

Plan: The conjugate base of a substance is simply the parent substance minus one proton, and the conjugate acid of a substance is the parent substance plus one proton.

Solve: (a) HClO_4 less one proton (H^+) is ClO_4^- . The other conjugate bases are HS^- , PH_3 , and CO_3^{2-} . (b) CN^- plus one proton (H^+) is HCN . The other conjugate acids are HSO_4^- , H_3O^+ , and H_2CO_3 .

Notice that the hydrogen carbonate ion (HCO_3^-) is amphiprotic. It can act as either an acid or a base.

PRACTICE EXERCISE

Write the formula for the conjugate acid of each of the following: HSO_3^- , F^- , PO_4^{3-} , CO .

Answers: H_2SO_3 , HF , HPO_4^{2-} , HCO^+


SAMPLE EXERCISE 16.2 | Writing Equations for Proton-Transfer Reactions

The hydrogen sulfite ion (HSO_3^-) is amphiprotic. (a) Write an equation for the reaction of HSO_3^- with water, in which the ion acts as an acid. (b) Write an equation for the reaction of HSO_3^- with water, in which the ion acts as a base. In both cases identify the conjugate acid–base pairs.


SOLUTION

Analyze and Plan: We are asked to write two equations representing reactions between HSO_3^- and water, one in which HSO_3^- should donate a proton to water, thereby acting as a Brønsted–Lowry acid, and one in which HSO_3^- should accept a proton from water, thereby acting as a base. We are also asked to identify the conjugate pairs in each equation.

Solve:



The conjugate pairs in this equation are HSO_3^- (acid) and SO_3^{2-} (conjugate base); and H_2O (base) and H_3O^+ (conjugate acid).



The conjugate pairs in this equation are H_2O (acid) and OH^- (conjugate base), and HSO_3^- (base) and H_2SO_3 (conjugate acid).

PRACTICE EXERCISE

When lithium oxide (Li_2O) is dissolved in water, the solution turns basic from the reaction of the oxide ion (O^{2-}) with water. Write the reaction that occurs, and identify the conjugate acid–base pairs.

Answer: $\text{O}^{2-}(aq) + \text{H}_2\text{O}(l) \longrightarrow \text{OH}^-(aq) + \text{OH}^-(aq)$. OH^- is the conjugate acid of the base O^{2-} . OH^- is also the conjugate base of the acid H_2O .

Relative Strengths of Acids and Bases

Some acids are better proton donors than others; likewise, some bases are better proton acceptors than others. If we arrange acids in order of their ability to donate a proton, we find that the more easily a substance gives up a proton, the less easily its conjugate base accepts a proton. Similarly, the more easily a base accepts a proton, the less easily its conjugate acid gives up a proton. In other words, *the stronger an acid, the weaker is its conjugate base; the stronger a base, the weaker is its conjugate acid*. Thus, if we know something about the strength of an acid (its ability to donate protons), we also know something about the strength of its conjugate base (its ability to accept protons).

The inverse relationship between the strengths of acids and the strengths of their conjugate bases is illustrated in Figure 16.4. Here we have grouped acids and bases into three broad categories based on their behavior in water.

1. A *strong acid* completely transfers its protons to water, leaving no undissociated molecules in solution. \rightleftharpoons (Section 4.3) Its conjugate base has a negligible tendency to be protonated (to abstract protons) in aqueous solution.
2. A *weak acid* only partially dissociates in aqueous solution and therefore exists in the solution as a mixture of acid molecules and their constituent ions. The conjugate base of a weak acid shows a slight ability to remove protons from water. (*The conjugate base of a weak acid is a weak base.*)
3. A substance with *negligible acidity*, such as CH_4 , contains hydrogen but does not demonstrate any acidic behavior in water. Its conjugate base is a strong base, reacting completely with water, abstracting protons to form OH^- ions.


	ACID	BASE		
100% ionized in H_2O	Strong	HCl	Cl^-	Negligible
		H_2SO_4	HSO_4^-	
		HNO_3	NO_3^-	
Acid strength increases ↑	Weak	$\text{H}_3\text{O}^+(aq)$	H_2O	Base strength increases ↓
		HSO_4^-	SO_4^{2-}	
		H_3PO_4	H_2PO_4^-	
		HF	F^-	
		CH_3COOH	CH_3COO^-	
		H_2CO_3	HCO_3^-	
		H_2S	HS^-	
		H_2PO_4^-	HPO_4^{2-}	
		NH_4^+	NH_3	
		HCO_3^-	CO_3^{2-}	
Negligible	Strong	H_2O	OH^-	100% protonated in H_2O
		OH^-	O^{2-}	
		H_2	H^-	
		CH_4	CH_3^-	

▲ **Figure 16.4** Relative strengths of some conjugate acid-base pairs. The two members of each pair are listed opposite each other in the two columns. The acids decrease in strength from top to bottom, whereas their conjugate bases increase in strength from top to bottom.


GIVE IT SOME THOUGHT

Using the three categories above, specify the strength of HNO_3 and the strength of its conjugate base, NO_3^- .

We can think of proton-transfer reactions as being governed by the relative abilities of two bases to abstract protons. For example, consider the proton transfer that occurs when an acid HX dissolves in water:




If H_2O (the base in the forward reaction) is a stronger base than X^- (the conjugate base of HX), then H_2O will abstract the proton from HX to produce H_3O^+ and X^- . As a result, the equilibrium will lie to the right. This describes the behavior of a strong acid in water. For example, when HCl dissolves in water, the solution consists almost entirely of H_3O^+ and Cl^- ions with a negligible concentration of HCl molecules.



H_2O is a stronger base than Cl^- (Figure 16.4), so H_2O acquires the proton to become the hydronium ion.

When X^- is a stronger base than H_2O , the equilibrium will lie to the left. This situation occurs when HX is a weak acid. For example, an aqueous solution of acetic acid (CH_3COOH) consists mainly of CH_3COOH molecules with only a relatively few H_3O^+ and CH_3COO^- ions.



CH_3COO^- is a stronger base than H_2O (Figure 16.4) and therefore abstracts the proton from H_3O^+ .

From these examples, we conclude that *in every acid–base reaction the position of the equilibrium favors transfer of the proton from the stronger acid to the stronger base to form the weaker acid and the weaker base*. As a result, the equilibrium mixture contains more of the weaker acid and weaker base and less of the stronger acid and stronger base.

SAMPLE EXERCISE 16.3 Predicting the Position of a Proton-Transfer Equilibrium

For the following proton-transfer reaction, use Figure 16.4 to predict whether the equilibrium lies predominantly to the left (that is, $K_c < 1$) or to the right ($K_c > 1$):




SOLUTION

Analyze: We are asked to predict whether the equilibrium shown lies to the right, favoring products, or to the left, favoring reactants.

Plan: This is a proton-transfer reaction, and the position of the equilibrium will favor the proton going to the stronger of two bases. The two bases in the equation are CO_3^{2-} , the base in the forward reaction as written, and SO_4^{2-} , the conjugate base of HSO_4^- . We can find the relative positions of these two bases in Figure 16.4 to determine which is the stronger base.



Solve: CO_3^{2-} appears lower in the right-hand column in Figure 16.4 and is therefore a stronger base than SO_4^{2-} . CO_3^{2-} , therefore, will get the proton preferentially to become HCO_3^- , while SO_4^{2-} will remain mostly unprotonated. The resulting equilibrium will lie to the right, favoring products (that is, $K_c > 1$).



Comment: Of the two acids in the equation, HSO_4^- and HCO_3^- , the stronger one gives up a proton more readily while the weaker one tends to retain its proton. Thus, the equilibrium favors the direction in which the proton moves from the stronger acid and becomes bonded to the stronger base.

PRACTICE EXERCISE


For each of the following reactions, use Figure 16.4 to predict whether the equilibrium lies predominantly to the left or to the right:



Answers: (a) left, (b) right

16.3 THE AUTOIONIZATION OF WATER

One of the most important chemical properties of water is its ability to act as either a Brønsted acid or a Brønsted base, depending on the circumstances. In the presence of an acid, water acts as a proton acceptor; in the presence of a base, water acts as a proton donor. In fact, one water molecule can donate a proton to another water molecule:



We call this process the **autoionization** of water. No individual molecule remains ionized for long; the reactions are extremely rapid in both directions.

At room temperature only about two out of every 10^9 molecules are ionized at any given instant. Thus, pure water consists almost entirely of H_2O molecules and is an extremely poor conductor of electricity. Nevertheless, the autoionization of water is very important, as we will soon see.

The Ion Product of Water


Because the autoionization of water (Equation 16.12) is an equilibrium process, we can write the following equilibrium-constant expression for it:

$$K_c = [\text{H}_3\text{O}^+][\text{OH}^-] \quad [16.13]$$

The term $[\text{H}_2\text{O}]$ is excluded from the equilibrium-constant expression because we exclude the concentrations of pure solids and liquids. ∞ (Section 15.4) Because this equilibrium-constant expression refers specifically to the autoionization of water, we use the symbol K_w to denote the equilibrium constant, which we call the **ion-product constant** for water. At 25°C , K_w equals 1.0×10^{-14} . Thus, we have

$$K_w = [\text{H}_3\text{O}^+][\text{OH}^-] = 1.0 \times 10^{-14} \quad (\text{at } 25^\circ\text{C}) \quad [16.14]$$

Because we use $\text{H}^+(\text{aq})$ and $\text{H}_3\text{O}^+(\text{aq})$ interchangeably to represent the hydrated proton, the autoionization reaction for water can also be written as



Likewise, the expression for K_w can be written in terms of either H_3O^+ or H^+ , and K_w has the same value in either case:

$$K_w = [\text{H}_3\text{O}^+][\text{OH}^-] = [\text{H}^+][\text{OH}^-] = 1.0 \times 10^{-14} \quad (\text{at } 25^\circ\text{C}) \quad [16.16]$$

This equilibrium-constant expression and the value of K_w at 25°C are extremely important, and you should commit them to memory.

What makes Equation 16.16 particularly useful is that it is applicable to pure water and to any aqueous solution. Although the equilibrium between $\text{H}^+(\text{aq})$ and $\text{OH}^-(\text{aq})$ as well as other ionic equilibria are affected somewhat by the presence of additional ions in solution, it is customary to ignore these ionic effects except in work requiring exceptional accuracy. Thus, Equation 16.16 is taken to be valid for any dilute aqueous solution, and it can be used to calculate either $[\text{H}^+]$ (if $[\text{OH}^-]$ is known) or $[\text{OH}^-]$ (if $[\text{H}^+]$ is known).

A solution in which $[\text{H}^+] = [\text{OH}^-]$ is said to be *neutral*. In most solutions H^+ and OH^- concentrations are not equal. As the concentration of one of these ions increases, the concentration of the other must decrease, so that the product of their concentrations equals 1.0×10^{-14} . In acidic solutions $[\text{H}^+]$ exceeds $[\text{OH}^-]$. In basic solutions $[\text{OH}^-]$ exceeds $[\text{H}^+]$.

SAMPLE EXERCISE 16.4 | Calculating $[\text{H}^+]$ for Pure Water

Calculate the values of $[\text{H}^+]$ and $[\text{OH}^-]$ in a neutral solution at 25°C .

SOLUTION

Analyze: We are asked to determine the concentrations of H^+ and OH^- ions in a neutral solution at 25°C .

Plan: We will use Equation 16.16 and the fact that, by definition, $[\text{H}^+] = [\text{OH}^-]$ in a neutral solution.

Solve: We will represent the concentration of $[\text{H}^+]$ and $[\text{OH}^-]$ in neutral solution with x . This gives

$$\begin{aligned} [\text{H}^+][\text{OH}^-] &= (x)(x) = 1.0 \times 10^{-14} \\ x^2 &= 1.0 \times 10^{-14} \\ x &= 1.0 \times 10^{-7} \text{ M} = [\text{H}^+] = [\text{OH}^-] \end{aligned}$$

In an acid solution $[\text{H}^+]$ is greater than $1.0 \times 10^{-7} \text{ M}$; in a basic solution $[\text{H}^+]$ is less than $1.0 \times 10^{-7} \text{ M}$.

PRACTICE EXERCISE

Indicate whether solutions with each of the following ion concentrations are neutral, acidic, or basic: (a) $[H^+] = 4 \times 10^{-9} M$; (b) $[OH^-] = 1 \times 10^{-7} M$; (c) $[OH^-] = 7 \times 10^{-13} M$.

Answers: (a) basic, (b) neutral, (c) acidic

SAMPLE EXERCISE 16.5 | Calculating $[H^+]$ from $[OH^-]$

Calculate the concentration of $H^+(aq)$ in (a) a solution in which $[OH^-]$ is $0.010 M$, (b) a solution in which $[OH^-]$ is $1.8 \times 10^{-9} M$. *Note:* In this problem and all that follow, we assume, unless stated otherwise, that the temperature is $25^\circ C$.

SOLUTION

Analyze: We are asked to calculate the hydronium ion concentration in an aqueous solution where the hydroxide concentration is known.

Plan: We can use the equilibrium-constant expression for the autoionization of water and the value of K_w to solve for each unknown concentration.

Solve:

(a) Using Equation 16.16, we have: $[H^+][OH^-] = 1.0 \times 10^{-14}$

$$[H^+] = \frac{(1.0 \times 10^{-14})}{[OH^-]} = \frac{1.0 \times 10^{-14}}{0.010} = 1.0 \times 10^{-12} M$$

This solution is basic because $[OH^-] > [H^+]$

(b) In this instance $[H^+] = \frac{(1.0 \times 10^{-14})}{[OH^-]} = \frac{1.0 \times 10^{-14}}{1.8 \times 10^{-9}} = 5.6 \times 10^{-6} M$

This solution is acidic because $[H^+] > [OH^-]$

PRACTICE EXERCISE

Calculate the concentration of $OH^-(aq)$ in a solution in which (a) $[H^+] = 2 \times 10^{-6} M$; (b) $[H^+] = [OH^-]$; (c) $[H^+] = 100 \times [OH^-]$. **Answers:** (a) $5 \times 10^{-9} M$, (b) $1.0 \times 10^{-7} M$, (c) $1.0 \times 10^{-8} M$

16.4 THE pH SCALE

The molar concentration of $H^+(aq)$ in an aqueous solution is usually very small. For convenience, we therefore usually express $[H^+]$ in terms of **pH**, which is the negative logarithm in base 10 of $[H^+]$.*

$$pH = -\log[H^+] \quad [16.17]$$

If you need to review the use of logs, see Appendix A.

We can use Equation 16.17 to calculate the pH of a neutral solution at $25^\circ C$ (that is, one in which $[H^+] = 1.0 \times 10^{-7} M$):

$$pH = -\log(1.0 \times 10^{-7}) = -(-7.00) = 7.00$$

The pH of a neutral solution is 7.00 at $25^\circ C$. Notice that the pH is reported with two decimal places. We do so because only the numbers to the right of the decimal point are the significant figures in a logarithm. Because our original value for the concentration ($1.0 \times 10^{-7} M$) has two significant figures, the corresponding pH has two decimal places (7.00).

What happens to the pH of a solution as we make the solution acidic? An acidic solution is one in which $[H^+] > 1.0 \times 10^{-7} M$. Because of the negative sign in Equation 16.17, *the pH decreases as $[H^+]$ increases*. For example, the pH of an acidic solution in which $[H^+] = 1.0 \times 10^{-3} M$ is

$$pH = -\log(1.0 \times 10^{-3}) = -(-3.00) = 3.00$$

At $25^\circ C$ the pH of an acidic solution is less than 7.00.

*Because $[H^+]$ and $[H_3O^+]$ are used interchangeably, you might see pH defined as $-\log[H_3O^+]$.

TABLE 16.1 Relationships among $[H^+]$, $[OH^-]$, and pH at 25 °C

Solution Type	$[H^+]$ (M)	$[OH^-]$ (M)	pH Value
Acidic	$>1.0 \times 10^{-7}$	$<1.0 \times 10^{-7}$	<7.00
Neutral	$=1.0 \times 10^{-7}$	$=1.0 \times 10^{-7}$	$=7.00$
Basic	$<1.0 \times 10^{-7}$	$>1.0 \times 10^{-7}$	>7.00

We can also calculate the pH of a basic solution, one in which $[OH^-] > 1.0 \times 10^{-7}$ M. Suppose $[OH^-] = 2.0 \times 10^{-3}$ M. We can use Equation 16.16 to calculate $[H^+]$ for this solution, and Equation 16.17 to calculate the pH:

$$[H^+] = \frac{K_w}{[OH^-]} = \frac{1.0 \times 10^{-14}}{2.0 \times 10^{-3}} = 5.0 \times 10^{-12} \text{ M}$$

$$\text{pH} = -\log(5.0 \times 10^{-12}) = 11.30$$

At 25 °C the pH of a basic solution is greater than 7.00. The relationships among $[H^+]$, $[OH^-]$, and pH are summarized in Table 16.1 and in Figure 16.5.

The pH values characteristic of several familiar solutions are shown in Figure 16.5. Notice that a change in $[H^+]$ by a factor of 10 causes the pH to change by 1. Thus, a solution of pH 6 has 10 times the concentration of $H^+(aq)$ as a solution of pH 7.



GIVE IT SOME THOUGHT

- (a) What is the significance of $\text{pH} = 7$? (b) How does the pH change as OH^- is added to the solution?

You might think that when $[H^+]$ is very small, as it is for some of the examples shown in Figure 16.5, it would be unimportant. Nothing is further from the truth. If $[H^+]$ is part of a kinetic rate law, then changing its concentration will

► Figure 16.5 H^+ concentrations and pH values of some common substances at 25 °C. The pH of a solution can be estimated using the benchmark concentrations of H^+ and OH^- corresponding to whole-number pH values.

	$[H^+]$ (M)	pH	pOH	$[OH^-]$ (M)
	$1 (1 \times 10^{-0})$	0.0	14.0	1×10^{-14}
More acidic ↑	1×10^{-1}	1.0	13.0	1×10^{-13}
Gastric juice	1×10^{-2}	2.0	12.0	1×10^{-12}
Lemon juice	1×10^{-3}	3.0	11.0	1×10^{-11}
Cola, vinegar	1×10^{-4}	4.0	10.0	1×10^{-10}
Wine	1×10^{-5}	5.0	9.0	1×10^{-9}
Tomatoes	1×10^{-6}	6.0	8.0	1×10^{-8}
Banana	1×10^{-7}	7.0	7.0	1×10^{-7}
Black coffee	1×10^{-8}	8.0	6.0	1×10^{-6}
Rain	1×10^{-9}	9.0	5.0	1×10^{-5}
Saliva	1×10^{-10}	10.0	4.0	1×10^{-4}
Milk	1×10^{-11}	11.0	3.0	1×10^{-3}
Human blood, tears	1×10^{-12}	12.0	2.0	1×10^{-2}
Egg white, seawater	1×10^{-13}	13.0	1.0	1×10^{-1}
Baking soda	1×10^{-14}	14.0	0.0	$1 (1 \times 10^{-0})$
Borax				
Milk of magnesia				
Lime water				
Household ammonia				
Household bleach				
NaOH, 0.1 M				
More basic ↓				

change the rate. ∞ (Section 14.3) Thus, if the rate law is first order in $[H^+]$ doubling its concentration will double the rate even if the change is merely from $1 \times 10^{-7} M$ to $2 \times 10^{-7} M$. In biological systems many reactions involve proton transfers and have rates that depend on $[H^+]$. Because the speeds of these reactions are crucial, the pH of biological fluids must be maintained within narrow limits. For example, human blood has a normal pH range of 7.35 to 7.45. Illness and even death can result if the pH varies much from this narrow range.

SAMPLE EXERCISE 16.6 | Calculating pH from $[H^+]$

Calculate the pH values for the two solutions described in Sample Exercise 16.5.

SOLUTION

Analyze: We are asked to determine the pH of aqueous solutions for which we have already calculated $[H^+]$.

Plan: We can calculate pH using its defining equation, Equation 16.17.

Solve:

(a) In the first instance we found $[H^+]$ to be $1.0 \times 10^{-12} M$.

$$\text{pH} = -\log(1.0 \times 10^{-12}) = -(-12.00) = 12.00$$

Because 1.0×10^{-12} has two significant figures, the pH has two decimal places, 12.00.

(b) For the second solution, $[H^+] = 5.6 \times 10^{-6} M$. Before performing the calculation, it is helpful to estimate the pH. To do so, we note that $[H^+]$ lies between 1×10^{-6} and 1×10^{-5} .

$$1 \times 10^{-6} < 5.6 \times 10^{-6} < 1 \times 10^{-5}$$

Thus, we expect the pH to lie between 6.0 and 5.0. We use Equation 16.17 to calculate the pH.

$$\text{pH} = -\log(5.6 \times 10^{-6}) = 5.25$$

Check: After calculating a pH, it is useful to compare it to your prior estimate. In this case the pH, as we predicted, falls between 6 and 5. Had the calculated pH and the estimate not agreed, we should have reconsidered our calculation or estimate or both.

PRACTICE EXERCISE

(a) In a sample of lemon juice $[H^+]$ is $3.8 \times 10^{-4} M$. What is the pH? (b) A commonly available window-cleaning solution has $[OH^-] = 1.9 \times 10^{-6} M$. What is the pH?

Answers: (a) 3.42, (b) $[H^+] = 5.3 \times 10^{-9} M$, so pH = 8.28

SAMPLE EXERCISE 16.7 | Calculating $[H^+]$ from pH

A sample of freshly pressed apple juice has a pH of 3.76. Calculate $[H^+]$.

SOLUTION

Analyze: We need to calculate $[H^+]$ from pH.

Plan: We will use Equation 16.17, $\text{pH} = -\log[H^+]$, for the calculation.

Solve: From Equation 16.17, we have

$$\text{pH} = -\log[H^+] = 3.76$$

Thus,

$$\log[H^+] = -3.76$$

To find $[H^+]$, we need to determine the *antilog* of -3.76 . Scientific calculators have an antilog function (sometimes labeled INV log or 10^x) that allows us to perform the calculation:

$$[H^+] = \text{antilog}(-3.76) = 10^{-3.76} = 1.7 \times 10^{-4} M$$

Comment: Consult the user's manual for your calculator to find out how to perform the antilog operation. The number of significant figures in $[H^+]$ is two because the number of decimal places in the pH is two.

Check: Because the pH is between 3.0 and 4.0, we know that $[H^+]$ will be between 1×10^{-3} and $1 \times 10^{-4} M$. Our calculated $[H^+]$ falls within this estimated range.

PRACTICE EXERCISE

A solution formed by dissolving an antacid tablet has a pH of 9.18. Calculate $[H^+]$.

Answer: $[H^+] = 6.6 \times 10^{-10} M$

pOH and Other “p” Scales

The negative log is also a convenient way of expressing the magnitudes of small quantities. We use the convention that the negative log of a quantity is labeled “p” (quantity). Thus, we can express the concentration of OH^- as pOH:

$$\text{pOH} = -\log[\text{OH}^-] \quad [16.18]$$

Likewise, $\text{p}K_w$ equals $-\log K_w$.

By taking the negative log of both sides of Equation 16.16,

$$-\log[\text{H}^+] + (-\log[\text{OH}^-]) = -\log K_w \quad [16.19]$$

we obtain the following useful expression:

$$\text{pH} + \text{pOH} = 14.00 \quad (\text{at } 25^\circ\text{C}) \quad [16.20]$$

We will see in Section 16.8 that p scales are also useful when working with other equilibrium constants.

GIVE IT SOME THOUGHT

If the pOH for a solution is 3.00, what is the pH of the solution? Is the solution acidic or basic?

Measuring pH

The pH of a solution can be measured quickly and accurately with a *pH meter* (Figure 16.6 ◀). A complete understanding of how this important device works requires a knowledge of electrochemistry, a subject we take up in Chapter 20. In brief, a pH meter consists of a pair of electrodes connected to a meter capable of measuring small voltages, on the order of millivolts. A voltage, which varies with the pH, is generated when the electrodes are placed in a solution. This voltage is read by the meter, which is calibrated to give pH.

The electrodes used with pH meters come in many shapes and sizes, depending on their intended use. Electrodes have even been developed that are so small that they can be inserted into single living cells to monitor the pH of the cell medium. Pocket-size pH meters are also available for use in environmental studies, in monitoring industrial effluents, and in agricultural work.

Although less precise, acid–base indicators can be used to measure pH. An acid–base indicator is a colored substance that itself can exist in either an acid or a base form. The two forms have different colors. Thus, the indicator turns one color in an acid and another color in a base. If you know the pH at which the indicator turns from one form to the other, you can determine whether a solution has a higher or lower pH than this value. Litmus, for example, changes color in the vicinity of pH 7. The color change, however, is not very sharp. Red litmus indicates a pH of about 5 or lower, and blue litmus indicates a pH of about 8 or higher.


Some of the more common indicators are listed in Figure 16.7 ▶. Methyl orange, for example, changes color over the pH interval from 3.1 to 4.4. Below pH 3.1 it is in the acid form, which is red. In the interval between 3.1 and 4.4, it is gradually converted to its basic form, which has a yellow color. By pH 4.4 the conversion is complete, and the solution is yellow. Paper tape that is impregnated with several indicators and comes complete with a comparator color scale is widely used for approximate determinations of pH.

GIVE IT SOME THOUGHT

If phenolphthalein turns pink when added to a solution, what can we conclude about the pH of the solution?



▲ **Figure 16.6** A digital pH meter. The device is a millivoltmeter, and the electrodes immersed in the solution being tested produce a voltage that depends on the pH of the solution.




▲ **Figure 16.7** Some common acid–base indicators. The pH ranges for the color changes of some common acid–base indicators. Most indicators have a useful range of about 2 pH units.

16.5 STRONG ACIDS AND BASES

The chemistry of an aqueous solution often depends critically on the pH of the solution. It is therefore important to examine how the pH of solutions relates to the concentrations of acids and bases. The simplest cases are those involving strong acids and strong bases. Strong acids and bases are *strong electrolytes*, existing in aqueous solution entirely as ions. There are relatively few common strong acids and bases, and we listed these substances in Table 4.2.

Strong Acids

The seven most common strong acids include six monoprotic acids (HCl, HBr, HI, HNO₃, HClO₃, and HClO₄), and one diprotic acid (H₂SO₄). Nitric acid (HNO₃) exemplifies the behavior of the monoprotic strong acids. For all practical purposes, an aqueous solution of HNO₃ consists entirely of H₃O⁺ and NO₃⁻ ions.



We have not used equilibrium arrows for Equation 16.21 because the reaction lies entirely to the right, the side with the ions. ∞ (Section 4.1) As noted in Section 16.3, we use H₃O⁺(aq) and H⁺(aq) interchangeably to represent the hydrated proton in water. Thus, we often simplify the equations for the ionization reactions of acids as follows:



In an aqueous solution of a strong acid, the acid is normally the only significant source of H⁺ ions.* As a result, calculating the pH of a solution of a strong monoprotic acid is straightforward because [H⁺] equals the original concentration of acid. In a 0.20 M solution of HNO₃(aq), for example, [H⁺] = [NO₃⁻] = 0.20 M. The situation with the diprotic acid H₂SO₄ is more complex, as we will see in Section 16.6.

*If the concentration of the acid is 10⁻⁶ M or less, we also need to consider H⁺ ions that result from the autoionization of H₂O. Normally, the concentration of H⁺ from H₂O is so small that it can be neglected.

SAMPLE EXERCISE 16.8 | Calculating the pH of a Strong Acid

What is the pH of a 0.040 M solution of HClO_4 ?

SOLUTION

Analyze and Plan: Because HClO_4 is a strong acid, it is completely ionized, giving $[\text{H}^+] = [\text{ClO}_4^-] = 0.040 \text{ M}$.

Solve: The pH of the solution is given by

$$\text{pH} = -\log(0.040) = 1.40.$$

Check: Because $[\text{H}^+]$ lies between 1×10^{-2} and 1×10^{-1} , the pH will be between 2.0 and 1.0. Our calculated pH falls within the estimated range. Furthermore, because the concentration has two significant figures, the pH has two decimal places.

PRACTICE EXERCISE

An aqueous solution of HNO_3 has a pH of 2.34. What is the concentration of the acid?

Answer: 0.0046 M

Strong Bases

There are relatively few common strong bases. The most common soluble strong bases are the ionic hydroxides of the alkali metals (group 1A) and the heavier alkaline earth metals (group 2A), such as NaOH , KOH , and Ca(OH)_2 . These compounds completely dissociate into ions in aqueous solution. Thus, a solution labeled 0.30 M NaOH consists of 0.30 M $\text{Na}^+(\text{aq})$ and 0.30 M $\text{OH}^-(\text{aq})$; there is essentially no undissociated NaOH .

SAMPLE EXERCISE 16.9 | Calculating the pH of a Strong Base

What is the pH of (a) a 0.028 M solution of NaOH , (b) a 0.0011 M solution of Ca(OH)_2 ?

SOLUTION

Analyze: We are asked to calculate the pH of two solutions of strong bases.

Plan: We can calculate each pH by either of two equivalent methods. First, we could use Equation 16.16 to calculate $[\text{H}^+]$ and then use Equation 16.17 to calculate the pH. Alternatively, we could use $[\text{OH}^-]$ to calculate pOH and then use Equation 16.20 to calculate the pH.

Solve:

(a) NaOH dissociates in water to give one OH^- ion per formula unit. Therefore, the OH^- concentration for the solution in (a) equals the stated concentration of NaOH , namely 0.028 M.

Method 1:

$$[\text{H}^+] = \frac{1.0 \times 10^{-14}}{0.028} = 3.57 \times 10^{-13} \text{ M} \quad \text{pH} = -\log(3.57 \times 10^{-13}) = 12.45$$

Method 2:

$$\text{pOH} = -\log(0.028) = 1.55 \quad \text{pH} = 14.00 - \text{pOH} = 12.45$$

(b) Ca(OH)_2 is a strong base that dissociates in water to give two OH^- ions per formula unit. Thus, the concentration of $\text{OH}^-(\text{aq})$ for the solution in part (b) is $2 \times (0.0011 \text{ M}) = 0.0022 \text{ M}$.

Method 1:

$$[\text{H}^+] = \frac{1.0 \times 10^{-14}}{0.0022} = 4.55 \times 10^{-12} \text{ M} \quad \text{pH} = -\log(4.55 \times 10^{-12}) = 11.34$$

Method 2:

$$\text{pOH} = -\log(0.0022) = 2.66 \quad \text{pH} = 14.00 - \text{pOH} = 11.34$$

PRACTICE EXERCISE

What is the concentration of a solution of (a) KOH for which the pH is 11.89; (b) Ca(OH)_2 for which the pH is 11.68?

Answers: (a) $7.8 \times 10^{-3} \text{ M}$, (b) $2.4 \times 10^{-3} \text{ M}$

Although all the hydroxides of the alkali metals (group 1A) are strong electrolytes, LiOH, RbOH, and CsOH are not commonly encountered in the laboratory. The hydroxides of the heavier alkaline earth metals, Ca(OH)₂, Sr(OH)₂, and Ba(OH)₂, are also strong electrolytes. They have limited solubilities, however, so they are used only when high solubility is not critical.

Strongly basic solutions are also created by certain substances that react with water to form OH⁻(aq). The most common of these contain the oxide ion. Ionic metal oxides, especially Na₂O and CaO, are often used in industry when a strong base is needed. The O²⁻ reacts with water to form OH⁻, leaving virtually no O²⁻ remaining in the solution:




Thus, a solution formed by dissolving 0.010 mol of Na₂O(s) in enough water to form 1.0 L of solution will have [OH⁻] = 0.020 M and a pH of 12.30.

GIVE IT SOME THOUGHT


The CH₃⁻ ion is the conjugate base of CH₄, and CH₄ shows no evidence of being an acid in water. What happens when CH₃⁻ is added to water?

16.6 WEAK ACIDS

Most acidic substances are weak acids and are therefore only partially ionized in aqueous solution. We can use the equilibrium constant for the ionization reaction to express the extent to which a weak acid ionizes. If we represent a general weak acid as HA, we can write the equation for its ionization reaction in either of the following ways, depending on whether the hydrated proton is represented as H₃O⁺(aq) or H⁺(aq):



or



Because H₂O is the solvent, it is omitted from the equilibrium-constant expression. ∞ (Section 15.4) Thus, we can write the equilibrium-constant expression as either

$$K_c = \frac{[\text{H}_3\text{O}^{+}][\text{A}^{-}]}{[\text{HA}]} \quad \text{or} \quad K_c = \frac{[\text{H}^{+}][\text{A}^{-}]}{[\text{HA}]}$$

As we did for the ion-product constant for the autoionization of water, we change the subscript on this equilibrium constant to indicate the type of equation to which it corresponds.




$$K_a = \frac{[\text{H}_3\text{O}^{+}][\text{A}^{-}]}{[\text{HA}]} \quad \text{or} \quad K_a = \frac{[\text{H}^{+}][\text{A}^{-}]}{[\text{HA}]} \quad [16.25]$$

The subscript *a* on *K_a* denotes that it is an equilibrium constant for the ionization of an acid, so *K_a* is called the **acid-dissociation constant**.

Table 16.2 shows the names, structures, and *K_a* values for several weak acids. Appendix D provides a more complete list. Many weak acids are organic compounds composed entirely of carbon, hydrogen, and oxygen. These compounds usually contain some hydrogen atoms bonded to carbon atoms and some bonded to oxygen atoms. In almost all cases the hydrogen atoms bonded to carbon do not ionize in water; instead, the acidic behavior of these compounds is due to the hydrogen atoms attached to oxygen atoms.

The magnitude of *K_a* indicates the tendency of the acid to ionize in water: *The larger the value of K_a, the stronger the acid.* Hydrofluoric acid (HF), for example, is the strongest acid listed in Table 16.2, and phenol (HOC₆H₅) is the weakest. Notice that *K_a* is typically less than 10⁻³.

TABLE 16.2 ■ Some Weak Acids in Water at 25 °C

Acid	Structural Formula*	Conjugate Base	Equilibrium Reaction	K_a
Hydrofluoric (HF)	$\text{H}-\text{F}$	F^-	$\text{HF}(aq) + \text{H}_2\text{O}(l) \rightleftharpoons \text{H}_3\text{O}^+(aq) + \text{F}^-(aq)$	6.8×10^{-4}
Nitrous (HNO_2)	$\text{H}-\text{O}-\text{N}=\text{O}$	NO_2^-	$\text{HNO}_2(aq) + \text{H}_2\text{O}(l) \rightleftharpoons \text{H}_3\text{O}^+(aq) + \text{NO}_2^-(aq)$	4.5×10^{-4}
Benzoic ($\text{C}_6\text{H}_5\text{COOH}$)		$\text{C}_6\text{H}_5\text{COO}^-$	$\text{C}_6\text{H}_5\text{COOH}(aq) + \text{H}_2\text{O}(l) \rightleftharpoons \text{H}_3\text{O}^+(aq) + \text{C}_6\text{H}_5\text{COO}^-(aq)$	6.3×10^{-5}
Acetic (CH_3COOH)		CH_3COO^-	$\text{CH}_3\text{COOH}(aq) + \text{H}_2\text{O}(l) \rightleftharpoons \text{H}_3\text{O}^+(aq) + \text{CH}_3\text{COO}^-(aq)$	1.8×10^{-5}
Hypochlorous (HClO)	$\text{H}-\text{O}-\text{Cl}$	ClO^-	$\text{HClO}(aq) + \text{H}_2\text{O}(l) \rightleftharpoons \text{H}_3\text{O}^+(aq) + \text{ClO}^-(aq)$	3.0×10^{-8}
Hydrocyanic (HCN)	$\text{H}-\text{C}\equiv\text{N}$	CN^-	$\text{HCN}(aq) + \text{H}_2\text{O}(l) \rightleftharpoons \text{H}_3\text{O}^+(aq) + \text{CN}^-(aq)$	4.9×10^{-10}
Phenol (HOC_6H_5)		$\text{C}_6\text{H}_5\text{O}^-$	$\text{HOC}_6\text{H}_5(aq) + \text{H}_2\text{O}(l) \rightleftharpoons \text{H}_3\text{O}^+(aq) + \text{C}_6\text{H}_5\text{O}^-(aq)$	1.3×10^{-10}

* The proton that ionizes is shown in blue.

Calculating K_a from pH

In order to calculate either the K_a value for a weak acid or the pH of its solutions, we will use many of the skills for solving equilibrium problems that we developed in Section 15.5. In many cases the small magnitude of K_a allows us to use approximations to simplify the problem. In doing these calculations, it is important to realize that proton-transfer reactions are generally very rapid. As a result, the measured or calculated pH for a weak acid always represents an equilibrium condition.

SAMPLE EXERCISE 16.10 | Calculating K_a from Measured pH


A student prepared a 0.10 M solution of formic acid (HCOOH) and measured its pH. The pH at 25 °C was found to be 2.38. Calculate K_a for formic acid at this temperature.

SOLUTION

Analyze: We are given the molar concentration of an aqueous solution of weak acid and the pH of the solution, and we are asked to determine the value of K_a for the acid.

Plan: Although we are dealing specifically with the ionization of a weak acid, this problem is very similar to the equilibrium problems we encountered in Chapter 15. We can solve this problem using the method first outlined in Sample Exercise 15.9, starting with the chemical reaction and a tabulation of initial and equilibrium concentrations.

Solve: The first step in solving any equilibrium problem is to write the equation for the equilibrium reaction. The ionization of formic acid can be written as follows:



The equilibrium-constant expression is

$$K_a = \frac{[\text{H}^+][\text{HCOO}^-]}{[\text{HCOOH}]}$$

From the measured pH, we can calculate $[\text{H}^+]$:

$$\begin{aligned} \text{pH} &= -\log[\text{H}^+] = 2.38 \\ \log[\text{H}^+] &= -2.38 \\ [\text{H}^+] &= 10^{-2.38} = 4.2 \times 10^{-3} \text{ M} \end{aligned}$$

We can do a little accounting to determine the concentrations of the species involved in the equilibrium. We imagine that the solution is initially 0.10 M in HCOOH molecules. We then consider the ionization of the acid into H^+ and HCOO^- . For each HCOOH molecule that ionizes, one H^+ ion and one HCOO^- ion are produced in solution. Because the pH measurement indicates that $[\text{H}^+] = 4.2 \times 10^{-3} \text{ M}$ at equilibrium, we can construct the following table:

$$\text{HCOOH}(aq) \rightleftharpoons \text{H}^+(aq) + \text{HCOO}^-(aq)$$

Initial	0.10 M	0	0
Change	$-4.2 \times 10^{-3} \text{ M}$	$+4.2 \times 10^{-3} \text{ M}$	$+4.2 \times 10^{-3} \text{ M}$
Equilibrium	$(0.10 - 4.2 \times 10^{-3}) \text{ M}$	$4.2 \times 10^{-3} \text{ M}$	$4.2 \times 10^{-3} \text{ M}$

Notice that we have neglected the very small concentration of $\text{H}^+(aq)$ that is due to the autoionization of H_2O . Notice also that the amount of HCOOH that ionizes is very small compared with the initial concentration of the acid. To the number of significant figures we are using, the subtraction yields 0.10 M:

$$(0.10 - 4.2 \times 10^{-3}) \text{ M} \approx 0.10 \text{ M}$$


We can now insert the equilibrium concentrations into the expression for K_a :

$$K_a = \frac{(4.2 \times 10^{-3})(4.2 \times 10^{-3})}{0.10} = 1.8 \times 10^{-4}$$

Check: The magnitude of our answer is reasonable because K_a for a weak acid is usually between 10^{-3} and 10^{-10} .

PRACTICE EXERCISE

Niacin, one of the B vitamins, has the following molecular structure:



A 0.020 M solution of niacin has a pH of 3.26. What is the acid-dissociation constant, K_a , for niacin?

Answers: 1.5×10^{-5}

Percent Ionization

We have seen that the magnitude of K_a indicates the strength of a weak acid. Another measure of acid strength is **percent ionization**, which is defined as

$$\text{Percent ionization} = \frac{\text{concentration ionized}}{\text{original concentration}} \times 100\% \quad [16.26]$$

The stronger the acid, the greater is the percent ionization.

For any acid, the concentration of acid that undergoes ionization equals the concentration of $\text{H}^+(aq)$ that forms, assuming that the autoionization of water is negligible. Thus, the percent ionization for an acid HA is also given by

$$\text{Percent ionization} = \frac{[\text{H}^+]_{\text{equilibrium}}}{[\text{HA}]_{\text{initial}}} \times 100\% \quad [16.27]$$

For example, a 0.035 M solution of HNO_2 contains $3.7 \times 10^{-3} \text{ M}$ $\text{H}^+(aq)$. Thus, the percent ionization is

$$\text{Percent ionization} = \frac{[\text{H}^+]_{\text{equilibrium}}}{[\text{HNO}_2]_{\text{initial}}} \times 100\% = \frac{3.7 \times 10^{-3} \text{ M}}{0.035 \text{ M}} \times 100\% = 11\%$$

SAMPLE EXERCISE 16.11 | Calculating Percent Ionization

A 0.10 M solution of formic acid (HCOOH) contains $4.2 \times 10^{-3} M H^+(aq)$. Calculate the percentage of the acid that is ionized.

SOLUTION

Analyze: We are given the molar concentration of an aqueous solution of weak acid and the equilibrium concentration of $H^+(aq)$ and asked to determine the percent ionization of the acid.

Plan: The percent ionization is given by Equation 16.27.

Solve:

$$\text{Percent ionization} = \frac{[H^+]_{\text{equilibrium}}}{[HCOOH]_{\text{initial}}} \times 100\% = \frac{4.2 \times 10^{-3} M}{0.10 M} \times 100\% = 4.2\%$$

PRACTICE EXERCISE


A 0.020 M solution of niacin has a pH of 3.26. Calculate the percent ionization of the niacin.

Answer: 2.7%

Using K_a to Calculate pH

Knowing the value of K_a and the initial concentration of the weak acid, we can calculate the concentration of $H^+(aq)$ in a solution of a weak acid. Let's calculate the pH of a 0.30 M solution of acetic acid (CH_3COOH), the weak acid responsible for the characteristic odor and acidity of vinegar, at 25 °C.

Our *first* step is to write the ionization equilibrium for acetic acid:



Notice that the hydrogen that ionizes is the one attached to an oxygen atom.

The *second* step is to write the equilibrium-constant expression and the value for the equilibrium constant. From Table 16.2, we have $K_a = 1.8 \times 10^{-5}$. Thus, we can write the following:

$$K_a = \frac{[H^+][CH_3COO^-]}{[CH_3COOH]} = 1.8 \times 10^{-5} \quad [16.29]$$

As the *third* step, we need to express the concentrations that are involved in the equilibrium reaction. This can be done with a little accounting, as described in Sample Exercise 16.10. Because we want to find the equilibrium value for $[H^+]$, let's call this quantity x . The concentration of acetic acid before any of it ionizes is 0.30 M. The chemical equation tells us that for each molecule of CH_3COOH that ionizes, one $H^+(aq)$ and one $CH_3COO^-(aq)$ are formed. Consequently, if x moles per liter of $H^+(aq)$ form at equilibrium, x moles per liter of $CH_3COO^-(aq)$ must also form, and x moles per liter of CH_3COOH must be ionized. This gives rise to the following table with the equilibrium concentrations shown on the last line:

	$CH_3COOH(aq)$	\rightleftharpoons	$H^+(aq)$	+	$CH_3COO^-(aq)$
Initial	0.30 M		0		0
Change	$-x M$		$+x M$		$+x M$
Equilibrium	$(0.30 - x) M$		$x M$		$x M$

As the *fourth* step of the problem, we need to substitute the equilibrium concentrations into the equilibrium-constant expression. The substitutions give the following equation:

$$K_a = \frac{[H^+][CH_3COO^-]}{[CH_3COOH]} = \frac{(x)(x)}{0.30 - x} = 1.8 \times 10^{-5} \quad [16.30]$$

This expression leads to a quadratic equation in x , which we can solve by using an equation-solving calculator or by using the quadratic formula. We can also simplify the problem, however, by noting that the value of K_a is quite small. As a result, we anticipate that the equilibrium will lie far to the left and that x will be very small compared to the initial concentration of acetic acid. Thus, we will *assume* that x is negligible compared to 0.30, so that $0.30 - x$ is essentially equal to 0.30.

$$0.30 - x \approx 0.30$$

As we will see, we can (and should!) check the validity of this assumption when we finish the problem. By using this assumption, Equation 16.30 now becomes

$$K_a = \frac{x^2}{0.30} = 1.8 \times 10^{-5}$$

Solving for x , we have

$$\begin{aligned} x^2 &= (0.30)(1.8 \times 10^{-5}) = 5.4 \times 10^{-6} \\ x &= \sqrt{5.4 \times 10^{-6}} = 2.3 \times 10^{-3} \\ [\text{H}^+] &= x = 2.3 \times 10^{-3} \text{ M} \\ \text{pH} &= -\log(2.3 \times 10^{-3}) = 2.64 \end{aligned}$$

We should now go back and check the validity of our simplifying assumption that $0.30 - x \approx 0.30$. The value of x we determined is so small that, for this number of significant figures, the assumption is entirely valid. We are thus satisfied that the assumption was a reasonable one to make. Because x represents the moles per liter of acetic acid that ionize, we see that, in this particular case, less than 1% of the acetic acid molecules ionize:

$$\text{Percent ionization of CH}_3\text{COOH} = \frac{0.0023 \text{ M}}{0.30 \text{ M}} \times 100\% = 0.77\%$$

As a general rule, if the quantity x is more than about 5% of the initial value, it is better to use the quadratic formula. You should always check the validity of any simplifying assumptions after you have finished solving a problem.

GIVE IT SOME THOUGHT

Why can we generally assume that the equilibrium concentration of a weak acid equals its initial concentration?

Finally, we can compare the pH value of this weak acid to a solution of a strong acid of the same concentration. The pH of the 0.30 M solution of acetic acid is 2.64. By comparison, the pH of a 0.30 M solution of a strong acid such as HCl is $-\log(0.30) = 0.52$. As expected, the pH of a solution of a weak acid is higher than that of a solution of a strong acid of the same molarity.

SAMPLE EXERCISE 16.12 | Using K_a to Calculate pH


Calculate the pH of a 0.20 M solution of HCN. (Refer to Table 16.2 or Appendix D for the value of K_a .)

SOLUTION

Analyze: We are given the molarity of a weak acid and are asked for the pH. From Table 16.2, K_a for HCN is 4.9×10^{-10} .

Plan: We proceed as in the example just worked in the text, writing the chemical equation and constructing a table of initial and equilibrium concentrations in which the equilibrium concentration of H^+ is our unknown.

Solve: Writing both the chemical equation for the ionization reaction that forms $\text{H}^+(\text{aq})$ and the equilibrium-constant (K_a) expression for the reaction:



Next, we tabulate the concentration of the species involved in the equilibrium reaction, letting $x = [\text{H}^+]$ at equilibrium:

	$\text{HCN}(aq)$	\rightleftharpoons	$\text{H}^+(aq)$	+	$\text{CN}^-(aq)$
Initial	0.20 M		0		0
Change	$-x$ M		$+x$ M		$+x$ M
Equilibrium	$(0.20 - x)$ M		x M		x M

Substituting the equilibrium concentrations from the table into the equilibrium-constant expression yields

$$K_a = \frac{(x)(x)}{0.20 - x} = 4.9 \times 10^{-10}$$

We next make the simplifying approximation that x , the amount of acid that dissociates, is small compared with the initial concentration of acid; that is,

$$0.20 - x \approx 0.20$$

Thus,

$$\frac{x^2}{0.20} = 4.9 \times 10^{-10}$$

Solving for x , we have

$$x^2 = (0.20)(4.9 \times 10^{-10}) = 0.98 \times 10^{-10}$$

$$x = \sqrt{0.98 \times 10^{-10}} = 9.9 \times 10^{-6} \text{ M} = [\text{H}^+]$$

A concentration of 9.9×10^{-6} M is much smaller than 5% of 0.20, the initial HCN concentration. Our simplifying approximation is therefore appropriate. We now calculate the pH of the solution:

$$\text{pH} = -\log[\text{H}^+] = -\log(9.9 \times 10^{-6}) = 5.00$$


PRACTICE EXERCISE


The K_a for niacin (Practice Exercise 16.10) is 1.5×10^{-5} . What is the pH of a 0.010 M solution of niacin?
Answer: 3.41

The properties of the acid solution that relate directly to the concentration of $\text{H}^+(aq)$, such as electrical conductivity and rate of reaction with an active metal, are much less evident for a solution of a weak acid than for a solution of a strong acid of the same concentration. Figure 16.8 \blacktriangledown presents an experiment that demonstrates this difference by comparing the behavior of 1 M CH_3COOH and 1 M HCl. The 1 M CH_3COOH contains only 0.004 M $\text{H}^+(aq)$, whereas the 1 M HCl solution contains 1 M $\text{H}^+(aq)$. As a result, the rate of reaction with the metal is much faster for the solution of HCl.

As the concentration of a weak acid increases, the equilibrium concentration of $\text{H}^+(aq)$ increases, as expected. However, as shown in Figure 16.9 \blacktriangleright , the percent ionization decreases as the concentration increases. Thus, the concentration of $\text{H}^+(aq)$ is not directly proportional to the concentration of the weak acid. For example, doubling the concentration of a weak acid does not double the concentration of $\text{H}^+(aq)$. This lack of proportionality between the concentration of a weak acid and the concentration of $\text{H}^+(aq)$ is demonstrated in Sample Exercise 16.13.

► Figure 16.8 Reaction rates for weak and strong acids. (a) The flask on the left contains 1 M CH_3COOH ; the one on the right contains 1 M HCl. Each balloon contains the same amount of magnesium metal. (b) When the Mg metal is dropped into the acid, H_2 gas is formed. The rate of H_2 formation is higher for the 1 M HCl solution on the right as evidenced by more gas in the balloon. Eventually, the same amount of H_2 forms in both cases.





◀ **Figure 16.9** The effect of concentration on ionization of a weak acid. The percent ionization of a weak acid decreases with increasing concentration. The data shown are for acetic acid.

■ SAMPLE EXERCISE 16.13 | Using K_a to Calculate Percent Ionization

Calculate the percentage of HF molecules ionized in (a) a 0.10 M HF solution, (b) a 0.010 M HF solution.

SOLUTION

Analyze: We are asked to calculate the percent ionization of two HF solutions of different concentration. From Appendix D, we find $K_a = 6.8 \times 10^{-4}$.

Plan: We approach this problem as we would previous equilibrium problems. We begin by writing the chemical equation for the equilibrium and tabulating the known and unknown concentrations of all species. We then substitute the equilibrium concentrations into the equilibrium-constant expression and solve for the unknown concentration, that of H^+ .

Solve:

(a) The equilibrium reaction and equilibrium concentrations are as follows:

	$HF(aq)$	\rightleftharpoons	$H^+(aq)$	+	$F^-(aq)$
Initial	0.10 M		0		0
Change	$-x$ M		$+x$ M		$+x$ M
Equilibrium	$(0.10 - x)$ M		x M		x M

The equilibrium-constant expression is

$$K_a = \frac{[H^+][F^-]}{[HF]} = \frac{(x)(x)}{0.10 - x} = 6.8 \times 10^{-4}$$

When we try solving this equation using the approximation $0.10 - x = 0.10$ (that is, by neglecting the concentration of acid that ionizes in comparison with the initial concentration), we obtain

$$x = 8.2 \times 10^{-3} M$$

Because this value is greater than 5% of 0.10 M, we should work the problem without the approximation, using an equation-solving calculator or the quadratic formula. Rearranging our equation and writing it in standard quadratic form, we have

$$\begin{aligned} x^2 &= (0.10 - x)(6.8 \times 10^{-4}) \\ &= 6.8 \times 10^{-5} - (6.8 \times 10^{-4})x \\ x^2 + (6.8 \times 10^{-4})x - 6.8 \times 10^{-5} &= 0 \end{aligned}$$

This equation can be solved using the standard quadratic formula.

$$x = \frac{-b \pm \sqrt{b^2 - 4ac}}{2a}$$

Substituting the appropriate numbers gives

$$\begin{aligned} x &= \frac{-6.8 \times 10^{-4} \pm \sqrt{(6.8 \times 10^{-4})^2 + 4(6.8 \times 10^{-5})}}{2} \\ &= \frac{-6.8 \times 10^{-4} \pm 1.6 \times 10^{-2}}{2} \end{aligned}$$

Of the two solutions, only the one that gives a positive value for x is chemically reasonable. Thus,

$$x = [\text{H}^+] = [\text{F}^-] = 7.9 \times 10^{-3} \text{ M}$$

From our result, we can calculate the percent of molecules ionized:

$$\begin{aligned} \text{Percent ionization of HF} &= \frac{\text{concentration ionized}}{\text{original concentration}} \times 100\% \\ &= \frac{7.9 \times 10^{-3} \text{ M}}{0.10 \text{ M}} \times 100\% = 7.9\% \end{aligned}$$

(b) Proceeding similarly for the 0.010 M solution, we have

$$\frac{x^2}{0.010 - x} = 6.8 \times 10^{-4}$$

Solving the resultant quadratic expression, we obtain

$$x = [\text{H}^+] = [\text{F}^-] = 2.3 \times 10^{-3} \text{ M}$$

The percentage of molecules ionized is

$$\frac{0.0023 \text{ M}}{0.010 \text{ M}} \times 100\% = 23\%$$

Comment: Notice that if we do not use the quadratic formula to solve the problem properly, we calculate 8.2% ionization for (a) and 26% ionization for (b). Notice also that in diluting the solution by a factor of 10, the percentage of molecules ionized increases by a factor of 3. This result is in accord with what we see in Figure 16.9. It is also what we would expect from Le Châtelier's principle. [Section 15.7](#) There are more "particles" or reaction components on the right side of the equation than on the left. Dilution causes the reaction to shift in the direction of the larger number of particles because this counters the effect of the decreasing concentration of particles.



PRACTICE EXERCISE

In Practice Exercise 16.11, we found that the percent ionization of niacin ($K_a = 1.5 \times 10^{-5}$) in a 0.020 M solution is 2.7%. Calculate the percentage of niacin molecules ionized in a solution that is (a) 0.010 M, (b) 1.0×10^{-3} M.

Answers: (a) 3.9%, (b) 12%

Polyprotic Acids

Many acids have more than one ionizable H atom. These acids are known as **polyprotic acids**. For example, each of the H atoms in sulfurous acid (H_2SO_3) can ionize in successive steps:



The acid-dissociation constants for these equilibria are labeled K_{a1} and K_{a2} . The numbers on the constants refer to the particular proton of the acid that is ionizing. Thus, K_{a2} always refers to the equilibrium involving removal of the second proton of a polyprotic acid.

In the preceding example K_{a2} is much smaller than K_{a1} . Because of electrostatic attractions, we would expect a positively charged proton to be lost more readily from the neutral H_2SO_3 molecule than from the negatively charged HSO_3^- ion. This observation is general: *It is always easier to remove the first proton from a polyprotic acid than to remove the second.* Similarly, for an acid with three ionizable protons, it is easier to remove the second proton than the third. Thus, the K_a values become successively smaller as successive protons are removed.





GIVE IT SOME THOUGHT

What is meant by the symbol K_{a3} for H_3PO_4 ?

The acid-dissociation constants for a few common polyprotic acids are listed in Table 16.3. Appendix D provides a more complete list. The structures for ascorbic and citric acids are shown in the margin. Notice that the K_a values for successive losses of protons from these acids usually differ by a factor of at least 10^3 . Notice also that the value of K_{a1} for sulfuric acid is listed simply as "large."

TABLE 16.3 Acid-Dissociation Constants of Some Common Polyprotic Acids

Name	Formula	K_{a1}	K_{a2}	K_{a3}
Ascorbic	$\text{H}_2\text{C}_6\text{H}_6\text{O}_6$	8.0×10^{-5}	1.6×10^{-12}	
Carbonic	H_2CO_3	4.3×10^{-7}	5.6×10^{-11}	
Citric	$\text{H}_3\text{C}_6\text{H}_5\text{O}_7$	7.4×10^{-4}	1.7×10^{-5}	4.0×10^{-7}
Oxalic	$\text{H}_2\text{C}_2\text{O}_4$	5.9×10^{-2}	6.4×10^{-5}	
Phosphoric	H_3PO_4	7.5×10^{-3}	6.2×10^{-8}	4.2×10^{-13}
Sulfurous	H_2SO_3	1.7×10^{-2}	6.4×10^{-8}	
Sulfuric	H_2SO_4	Large	1.2×10^{-2}	
Tartaric	$\text{H}_2\text{C}_4\text{H}_4\text{O}_6$	1.0×10^{-3}	4.6×10^{-5}	



Sulfuric acid is a strong acid with respect to the removal of the first proton. Thus, the reaction for the first ionization step lies completely to the right:




HSO_4^- , on the other hand, is a weak acid for which $K_{a2} = 1.2 \times 10^{-2}$.

Because K_{a1} is so much larger than subsequent dissociation constants for these polyprotic acids, most of the $\text{H}^+(aq)$ in the solution comes from the first ionization reaction. As long as successive K_a values differ by a factor of 10^3 or more, it is possible to obtain a satisfactory estimate of the pH of polyprotic acid solutions by treating them as if they were monoprotic acids, considering only K_{a1} .

SAMPLE EXERCISE 16.14 | Calculating the pH of a Polyprotic Acid Solution

The solubility of CO_2 in pure water at 25°C and 0.1 atm pressure is 0.0037 M. The common practice is to assume that all of the dissolved CO_2 is in the form of carbonic acid (H_2CO_3), which is produced by reaction between the CO_2 and H_2O :



What is the pH of a 0.0037 M solution of H_2CO_3 ?

SOLUTION

Analyze: We are asked to determine the pH of a 0.0037 M solution of a polyprotic acid.

Plan: H_2CO_3 is a diprotic acid; the two acid-dissociation constants, K_{a1} and K_{a2} (Table 16.3), differ by more than a factor of 10^3 . Consequently, the pH can be determined by considering only K_{a1} , thereby treating the acid as if it were a monoprotic acid.

Solve: Proceeding as in Sample Exercises 16.12 and 16.13, we can write the equilibrium reaction and equilibrium concentrations as follows:

$$\text{H}_2\text{CO}_3(aq) \rightleftharpoons \text{H}^+(aq) + \text{HCO}_3^-(aq)$$

Initial	0.0037 M	0	0
Change	$-x$ M	$+x$ M	$+x$ M
Equilibrium	$(0.0037 - x)$ M	x M	x M

The equilibrium-constant expression is as follows:

$$K_{a1} = \frac{[\text{H}^+][\text{HCO}_3^-]}{[\text{H}_2\text{CO}_3]} = \frac{(x)(x)}{0.0037 - x} = 4.3 \times 10^{-7}$$

Solving this equation using an equation-solving calculator, we get

$$x = 4.0 \times 10^{-5} \text{ M}$$

Alternatively, because K_{a1} is small, we can make the simplifying approximation that x is small, so that

$$0.0037 - x \approx 0.0037$$

Thus,

$$\frac{(x)(x)}{0.0037} = 4.3 \times 10^{-7}$$

Solving for x , we have

$$x^2 = (0.0037)(4.3 \times 10^{-7}) = 1.6 \times 10^{-9}$$

$$x = [\text{H}^+] = [\text{HCO}_3^-] = \sqrt{1.6 \times 10^{-9}} = 4.0 \times 10^{-5} \text{ M}$$

The small value of x indicates that our simplifying assumption was justified. The pH is therefore

$$\text{pH} = -\log[\text{H}^+] = -\log(4.0 \times 10^{-5}) = 4.40$$

Comment: If we were asked to solve for $[\text{CO}_3^{2-}]$, we would need to use K_{a2} . Let's illustrate that calculation. Using the values of $[\text{HCO}_3^-]$ and $[\text{H}^+]$ calculated above, and setting $[\text{CO}_3^{2-}] = y$, we have the following initial and equilibrium concentration values:

	$\text{HCO}_3^-(aq)$	\rightleftharpoons	$\text{H}^+(aq)$	+	$\text{CO}_3^{2-}(aq)$
Initial	$4.0 \times 10^{-5} M$		$4.0 \times 10^{-5} M$		0
Change	$-y M$		$+y M$		$+y M$
Equilibrium	$(4.0 \times 10^{-5} - y) M$		$(4.0 \times 10^{-5} + y) M$		$y M$

Assuming that y is small compared to 4.0×10^{-5} , we have

$$K_{a2} = \frac{[\text{H}^+][\text{CO}_3^{2-}]}{[\text{HCO}_3^-]} = \frac{(4.0 \times 10^{-5})(y)}{4.0 \times 10^{-5}} = 5.6 \times 10^{-11}$$

$$y = 5.6 \times 10^{-11} M = [\text{CO}_3^{2-}]$$

The value calculated for y is indeed very small compared to 4.0×10^{-5} , showing that our assumption was justified. It also shows that the ionization of HCO_3^- is negligible compared to that of H_2CO_3 , as far as production of H^+ is concerned. However, it is the *only* source of CO_3^{2-} , which has a very low concentration in the solution. Our calculations thus tell us that in a solution of carbon dioxide in water, most of the CO_2 is in the form of CO_2 or H_2CO_3 , a small fraction ionizes to form H^+ and HCO_3^- , and an even smaller fraction ionizes to give CO_3^{2-} . Notice also that $[\text{CO}_3^{2-}]$ is numerically equal to K_{a2} .


■ PRACTICE EXERCISE

(a) Calculate the pH of a 0.020 M solution of oxalic acid ($\text{H}_2\text{C}_2\text{O}_4$). (See Table 16.3 for K_{a1} and K_{a2} .) (b) Calculate the concentration of oxalate ion, $[\text{C}_2\text{O}_4^{2-}]$, in this solution.

Answers: (a) pH = 1.80, (b) $[\text{C}_2\text{O}_4^{2-}] = 6.4 \times 10^{-5} M$

16.7 WEAK BASES


Many substances behave as weak bases in water. Weak bases react with water, abstracting protons from H_2O , thereby forming the conjugate acid of the base and OH^- ions.



The equilibrium-constant expression for this reaction can be written as

$$K_b = \frac{[\text{BH}^+][\text{OH}^-]}{[\text{B}]} \quad [16.34]$$

Water is the solvent, so it is omitted from the equilibrium-constant expression. The most commonly encountered weak base is ammonia.



As with K_w and K_a , the subscript "b" denotes that this equilibrium constant refers to a particular type of reaction, namely the ionization of a weak base in water. The constant K_b is called the **base-dissociation constant**. The constant K_b always refers to the equilibrium in which a base reacts with H_2O to form the corresponding conjugate acid and OH^- .

Table 16.4 lists the names, formulas, Lewis structures, equilibrium reactions, and values of K_b for several weak bases in water. Appendix D includes a more extensive list. These bases contain one or more lone pairs of electrons because a lone pair is necessary to form the bond with H^+ . Notice that in the neutral molecules in Table 16.4, the lone pairs are on nitrogen atoms. The other bases listed are anions derived from weak acids.

TABLE 16.4 ■ Some Weak Bases and Their Aqueous Solution Equilibria

Base	Lewis Structure	Conjugate Acid	Equilibrium Reaction	K_b
Ammonia (NH ₃)		NH ₄ ⁺	NH ₃ + H ₂ O ⇌ NH ₄ ⁺ + OH ⁻	1.8 × 10 ⁻⁵
Pyridine (C ₅ H ₅ N)		C ₅ H ₅ NH ⁺	C ₅ H ₅ N + H ₂ O ⇌ C ₅ H ₅ NH ⁺ + OH ⁻	1.7 × 10 ⁻⁹
Hydroxylamine (H ₂ NOH)		H ₃ NOH ⁺	H ₂ NOH + H ₂ O ⇌ H ₃ NOH ⁺ + OH ⁻	1.1 × 10 ⁻⁸
Methylamine (NH ₂ CH ₃)		NH ₃ CH ₃ ⁺	NH ₂ CH ₃ + H ₂ O ⇌ NH ₃ CH ₃ ⁺ + OH ⁻	4.4 × 10 ⁻⁴
Hydrosulfide ion (HS ⁻)		H ₂ S	HS ⁻ + H ₂ O ⇌ H ₂ S + OH ⁻	1.8 × 10 ⁻⁷
Carbonate ion (CO ₃ ²⁻)		HCO ₃ ⁻	CO ₃ ²⁻ + H ₂ O ⇌ HCO ₃ ⁻ + OH ⁻	1.8 × 10 ⁻⁴
Hypochlorite ion (ClO ⁻)		HClO	ClO ⁻ + H ₂ O ⇌ HClO + OH ⁻	3.3 × 10 ⁻⁷

SAMPLE EXERCISE 16.15 | Using K_b to Calculate OH⁻


Calculate the concentration of OH⁻ in a 0.15 M solution of NH₃.

SOLUTION

Analyze: We are given the concentration of a weak base and are asked to determine the concentration of OH⁻.

Plan: We will use essentially the same procedure here as used in solving problems involving the ionization of weak acids; that is, we write the chemical equation and tabulate initial and equilibrium concentrations.

Solve: We first write the ionization reaction and the corresponding equilibrium-constant (K_b) expression:



$$K_b = \frac{[\text{NH}_4^+][\text{OH}^-]}{[\text{NH}_3]} = 1.8 \times 10^{-5}$$

We then tabulate the equilibrium concentrations involved in the equilibrium:

	$\text{NH}_3(aq) + \text{H}_2\text{O}(l) \rightleftharpoons \text{NH}_4^+(aq) + \text{OH}^-(aq)$			
Initial	0.15 M	—	0	0
Change	-x M	—	+x M	+x M
Equilibrium	(0.15 - x) M	—	x M	x M

(We ignore the concentration of H₂O because it is not involved in the equilibrium-constant expression.) Inserting these quantities into the equilibrium-constant expression gives the following:

$$K_b = \frac{[\text{NH}_4^+][\text{OH}^-]}{[\text{NH}_3]} = \frac{(x)(x)}{0.15 - x} = 1.8 \times 10^{-5}$$

Because K_b is small, we can neglect the small amount of NH₃ that reacts with water, as compared to the total NH₃ concentration; that is, we can neglect x relative to 0.15 M. Then we have

$$\begin{aligned} \frac{x^2}{0.15} &= 1.8 \times 10^{-5} \\ x^2 &= (0.15)(1.8 \times 10^{-5}) = 2.7 \times 10^{-6} \\ x &= [\text{NH}_4^+] = [\text{OH}^-] = \sqrt{2.7 \times 10^{-6}} = 1.6 \times 10^{-3} M \end{aligned}$$

Check: The value obtained for x is only about 1% of the NH_3 concentration, 0.15 M. Therefore, neglecting x relative to 0.15 was justified.


Comment: You may be asked to find the pH of a solution of a weak base. Once you have found $[\text{OH}^-]$, you can proceed as in Sample Exercise 16.9, where we calculated the pH of a strong base. In the present sample exercise, we have seen that the 0.15 M solution of NH_3 contains $[\text{OH}^-] = 1.6 \times 10^{-3}$ M. Thus, $\text{pOH} = -\log(1.6 \times 10^{-3}) = 2.80$, and $\text{pH} = 14.00 - 2.80 = 11.20$. The pH of the solution is above 7 because we are dealing with a solution of a base.

■ PRACTICE EXERCISE

Which of the following compounds should produce the highest pH as a 0.05 M solution: pyridine, methylamine, or nitrous acid?
Answer: methylamine (because it has the largest K_b value of the two amine bases in the list)


Types of Weak Bases

How can we recognize from a chemical formula whether a molecule or ion is able to behave as a weak base? Weak bases fall into two general categories. The first category contains neutral substances that have an atom with a nonbonding pair of electrons that can serve as a proton acceptor. Most of these bases, including all of the uncharged bases listed in Table 16.4, contain a nitrogen atom. These substances include ammonia and a related class of compounds called **amines**. In organic amines, one or more of the N—H bonds in NH_3 is replaced with a bond between N and C. Thus, the replacement of one N—H bond in NH_3 with a N—CH₃ bond gives methylamine, NH_2CH_3 (usually written CH_3NH_2). Like NH_3 , amines can abstract a proton from a water molecule by forming an additional N—H bond, as shown here for methylamine:



The chemical formula for the conjugate acid of methylamine is usually written CH_3NH_3^+ .

The second general category of weak bases consists of the anions of weak acids. In an aqueous solution of sodium hypochlorite (NaClO), for example, NaClO dissociates to give Na^+ and ClO^- ions. The Na^+ ion is always a spectator ion in acid-base reactions. \rightleftharpoons (Section 4.3) The ClO^- ion, however, is the conjugate base of a weak acid, hypochlorous acid. Consequently, the ClO^- ion acts as a weak base in water:



■ SAMPLE EXERCISE 16.16 | Using pH to Determine the Concentration of a Salt

A solution made by adding solid sodium hypochlorite (NaClO) to enough water to make 2.00 L of solution has a pH of 10.50. Using the information in Equation 16.37, calculate the number of moles of NaClO that were added to the water.

SOLUTION

Analyze: We are given the pH of a 2.00-L solution of NaClO and must calculate the number of moles of NaClO needed to raise the pH to 10.50. NaClO is an ionic compound consisting of Na^+ and ClO^- ions. As such, it is a strong electrolyte that completely dissociates in solution into Na^+ , which is a spectator ion, and ClO^- ion, which is a weak base with $K_b = 3.3 \times 10^{-7}$ (Equation 16.37).

Plan: From the pH, we can determine the equilibrium concentration of OH^- . We can then construct a table of initial and equilibrium concentrations in which the initial concentration of ClO^- is our unknown. We can calculate $[\text{ClO}^-]$ using the equilibrium-constant expression, K_b .

Solve: We can calculate $[\text{OH}^-]$ by using either Equation 16.16 or Equation 16.20; we will use the latter method here:

$$\begin{aligned} \text{pOH} &= 14.00 - \text{pH} = 14.00 - 10.50 = 3.50 \\ [\text{OH}^-] &= 10^{-3.50} = 3.2 \times 10^{-4} \text{ M} \end{aligned}$$

This concentration is high enough that we can assume that Equation 16.37 is the only source of OH^- ; that is, we can neglect any OH^- produced by the autoionization of H_2O . We now assume a value of x for the initial concentration of ClO^- and solve the equilibrium problem in the usual way.

$$\text{ClO}^-(aq) + \text{H}_2\text{O}(l) \rightleftharpoons \text{HClO}(aq) + \text{OH}^-(aq)$$

Initial	$x \text{ M}$	—	0	0
Change	$-3.2 \times 10^{-4} \text{ M}$	—	$+3.2 \times 10^{-4} \text{ M}$	$+3.2 \times 10^{-4} \text{ M}$
Equilibrium	$(x - 3.2 \times 10^{-4}) \text{ M}$	—	$3.2 \times 10^{-4} \text{ M}$	$3.2 \times 10^{-4} \text{ M}$

We now use the expression for the base-dissociation constant to solve for x :

$$K_b = \frac{[\text{HClO}][\text{OH}^-]}{[\text{ClO}^-]} = \frac{(3.2 \times 10^{-4})^2}{x - 3.2 \times 10^{-4}} = 3.3 \times 10^{-7}$$

Thus

$$x = \frac{(3.2 \times 10^{-4})^2}{3.3 \times 10^{-7}} + (3.2 \times 10^{-4}) = 0.31 \text{ M}$$

We say that the solution is 0.31 M in NaClO even though some of the ClO^- ions have reacted with water. Because the solution is 0.31 M in NaClO and the total volume of solution is 2.00 L, 0.62 mol of NaClO is the amount of the salt that was added to the water.



■ PRACTICE EXERCISE

A solution of NH_3 in water has a pH of 11.17. What is the molarity of the solution?

Answer: 0.12 M

16.8 RELATIONSHIP BETWEEN K_a AND K_b

We have seen in a qualitative way that the stronger acids have the weaker conjugate bases. To see if we can find a corresponding *quantitative* relationship, let's consider the NH_4^+ and NH_3 conjugate acid–base pair. Each of these species reacts with water:




Each of these equilibria is expressed by a characteristic dissociation constant:

$$K_a = \frac{[\text{NH}_3][\text{H}^+]}{[\text{NH}_4^+]}$$

$$K_b = \frac{[\text{NH}_4^+][\text{OH}^-]}{[\text{NH}_3]}$$

When Equations 16.38 and 16.39 are added together, the NH_4^+ and NH_3 species cancel and we are left with just the autoionization of water.



Recall that when two equations are added to give a third, the equilibrium constant associated with the third equation equals the product of the equilibrium constants for the two equations added together. ∞ (Section 15.3)

Applying this rule to our present example, when we multiply K_a and K_b , we obtain the following:

$$\begin{aligned} K_a \times K_b &= \left(\frac{[\text{NH}_3][\text{H}^+]}{[\text{NH}_4^+]} \right) \left(\frac{[\text{NH}_4^+][\text{OH}^-]}{[\text{NH}_3]} \right) \\ &= [\text{H}^+][\text{OH}^-] = K_w \end{aligned}$$

By using Equation 16.40, we can calculate K_b for any weak base if we know K_a for its conjugate acid. Similarly, we can calculate K_a for a weak acid if we know K_b for its conjugate base. As a practical consequence, ionization constants are often listed for only one member of a conjugate acid–base pair. For example, Appendix D does not contain K_b values for the anions of weak acids because they can be readily calculated from the tabulated K_a values for their conjugate acids.

If you look up the values for acid- or base-dissociation constants in a chemistry handbook, you may find them expressed as pK_a or pK_b (that is, as $-\log K_a$ or $-\log K_b$). ∞ (Section 16.4) Equation 16.40 can be written in terms of pK_a and pK_b by taking the negative log of both sides:

$$pK_a + pK_b = pK_w = 14.00 \quad \text{at } 25^\circ\text{C} \quad [16.41]$$

SAMPLE EXERCISE 16.17 | Calculating K_a or K_b for a Conjugate Acid–Base Pair

Calculate (a) the base-dissociation constant, K_b , for the fluoride ion (F^-); (b) the acid-dissociation constant, K_a , for the ammonium ion (NH_4^+).

SOLUTION

Analyze: We are asked to determine dissociation constants for F^- , the conjugate base of HF, and NH_4^+ , the conjugate acid of NH_3 .

Plan: Although neither F^- nor NH_4^+ appears in the tables, we can find the tabulated values for ionization constants for HF and NH_3 , and use the relationship between K_a and K_b to calculate the ionization constants for each of the conjugates.

Solve:

(a) K_a for the weak acid, HF, is given in Table 16.2 and Appendix D as $K_a = 6.8 \times 10^{-4}$. We can use Equation 16.40 to calculate K_b for the conjugate base, F^- :


$$K_b = \frac{K_w}{K_a} = \frac{1.0 \times 10^{-14}}{6.8 \times 10^{-4}} = 1.5 \times 10^{-11}$$

(b) K_b for NH_3 is listed in Table 16.4 and in Appendix D as $K_b = 1.8 \times 10^{-5}$. Using Equation 16.40, we can calculate K_a for the conjugate acid, NH_4^+ :

$$K_a = \frac{K_w}{K_b} = \frac{1.0 \times 10^{-14}}{1.8 \times 10^{-5}} = 5.6 \times 10^{-10}$$

PRACTICE EXERCISE

(a) Which of the following anions has the largest base-dissociation constant: NO_2^- , PO_4^{3-} , or N_3^- ? (b) The base quinoline has the following structure:



Its conjugate acid is listed in handbooks as having a pK_a of 4.90. What is the base-dissociation constant for quinoline?

Answers: (a) PO_4^{3-} ($K_b = 2.4 \times 10^{-2}$), (b) 7.9×10^{-10}

16.9 ACID–BASE PROPERTIES OF SALT SOLUTIONS

Even before you began this chapter, you were undoubtedly aware of many substances that are acidic, such as HNO_3 , HCl , and H_2SO_4 , and others that are basic, such as $NaOH$ and NH_3 . However, our recent discussions have indicated that ions can also exhibit acidic or basic properties. For example, we calculated K_a for NH_4^+ and K_b for F^- in Sample Exercise 16.17. Such behavior implies that salt solutions can be acidic or basic. Before proceeding with further discussions of acids and bases, let's examine the way dissolved salts can affect pH.

Because nearly all salts are strong electrolytes, we can assume that when salts dissolve in water, they are completely dissociated. Consequently, the acid-base properties of salt solutions are due to the behavior of their constituent cations and anions. Many ions are able to react with water to generate $\text{H}^+(\text{aq})$ or $\text{OH}^-(\text{aq})$. This type of reaction is often called **hydrolysis**. The pH of an aqueous salt solution can be predicted qualitatively by considering the ions of which the salt is composed.


An Anion's Ability to React with Water

In general, an anion, X^- , in solution can be considered the conjugate base of an acid. For example, Cl^- is the conjugate base of HCl , and CH_3COO^- is the conjugate base of CH_3COOH . Whether an anion reacts with water to produce hydroxide depends upon the strength of the acid to which it is conjugate. To identify the acid and assess its strength, we can simply add a proton to the anion's formula:



If the acid determined in this way is one of the strong acids listed at the beginning of Section 16.5, then the anion in question will have a negligible tendency to abstract protons from water. ∞ (Section 16.2) Consequently, the anion X^- will not affect the pH of the solution. The presence of Cl^- in an aqueous solution, for example, does not result in the production of any OH^- and does not affect the pH. Thus, Cl^- is always a spectator in acid-base chemistry.

Conversely, if HX is *not* one of the seven strong acids, then it is a weak acid. In this case the conjugate base X^- is a weak base. This anion will therefore react to a small extent with water to produce the weak acid and hydroxide ions:



The OH^- ion generated in this way increases the pH of the solution, making it basic. Acetate ion (CH_3COO^-), for example, being the conjugate base of a weak acid, reacts with water to produce acetic acid and hydroxide ions, thereby increasing the pH of the solution.*




GIVE IT SOME THOUGHT

What effect will each of the following ions have on the pH of a solution: NO_3^- and CO_3^{2-} ?

Anions that still have ionizable protons, such as HSO_3^- , are amphiprotic. ∞ (Section 16.2) They can act as either acids or bases. Their behavior toward water will be determined by the relative magnitudes of K_a and K_b for the ion, as shown in Sample Exercise 16.19. If $K_a > K_b$, the ion will cause the solution to be acidic. If $K_b > K_a$, the solution will be basic.

A Cation's Ability to React with Water

Polyatomic cations whose formulas contain one or more protons can be considered the conjugate acids of weak bases. NH_4^+ , for example, is the conjugate acid of the weak base NH_3 . Thus, NH_4^+ is a weak acid and will donate a proton to water, producing hydronium ions and thereby lowering the pH:



*These rules apply to what are called *normal salts*. These salts contain no ionizable protons on the anion. The pH of an acid salt (such as NaHCO_3 or NaH_2PO_4) is affected by the hydrolysis of the anion and by its acid dissociation, as shown in Sample Exercise 16.19.

Most metal ions can also react with water to decrease the pH of an aqueous solution. The mechanism by which metal ions produce acidic solutions is described in Section 16.11. However, ions of alkali metals and of the heavier alkaline earth metals do not react with water and therefore do not affect pH. Note that these exceptions are the cations found in the strong bases. ∞ (Section 16.5)

GIVE IT SOME THOUGHT

Which of the following cations has no effect on the pH of a solution: K^+ , Fe^{2+} , or Al^{3+} ?


Combined Effect of Cation and Anion in Solution

If an aqueous salt solution contains an anion that does not react with water and a cation that does not react with water, we expect the pH to be neutral. If the solution contains an anion that reacts with water to produce hydroxide and a cation that does not react with water, we expect the pH to be basic. If the solution contains a cation that reacts with water to produce hydronium and an anion that does not react with water, we expect the pH to be acidic. Finally, a solution may contain an anion and a cation *both* capable of reacting with water. In this case both hydroxide and hydronium will be produced. Whether the solution is basic, neutral, or acidic will depend upon the relative abilities of the ions to react with water.

To summarize:

1. An anion that is the conjugate base of a strong acid, for example, Br^- , will not affect the pH of a solution. (It will be a spectator ion in acid–base chemistry.)
2. An anion that is the conjugate base of a weak acid, for example, CN^- , will cause an increase in pH.
3. A cation that is the conjugate acid of a weak base, for example, $CH_3NH_3^+$, will cause a decrease in pH.
4. The cations of group 1A and heavier members of group 2A (Ca^{2+} , Sr^{2+} , and Ba^{2+}) will not affect pH. These are the cations of the strong Arrhenius bases. (They will be spectator ions in acid–base chemistry.)
5. Other metal ions will cause a decrease in pH.
6. When a solution contains both the conjugate base of a weak acid and the conjugate acid of a weak base, the ion with the larger equilibrium constant, K_a or K_b , will have the greater influence on the pH.

Figure 16.11 \blacktriangledown demonstrates the influence of several salts on pH.



\blacktriangleleft **Figure 16.11 Salt solutions can be neutral, acidic, or basic.** These three solutions contain the acid–base indicator bromthymol blue. (a) The NaCl solution is neutral (pH = 7.0); (b) the NH_4Cl solution is acidic (pH = 3.5); (c) the NaClO solution is basic (pH = 9.5).

SAMPLE EXERCISE 16.18 | Determining Whether Salt Solutions Are Acidic, Basic, or Neutral

Determine whether aqueous solutions of each of the following salts will be acidic, basic, or neutral: (a) $\text{Ba}(\text{CH}_3\text{COO})_2$, (b) NH_4Cl , (c) $\text{CH}_3\text{NH}_3\text{Br}$, (d) KNO_3 , (e) $\text{Al}(\text{ClO}_4)_3$.

SOLUTION

Analyze: We are given the chemical formulas of five ionic compounds (salts) and asked whether their aqueous solutions will be acidic, basic, or neutral.

Plan: We can determine whether a solution of a salt is acidic, basic, or neutral by identifying the ions in solution and by assessing how each ion will affect the pH.

Solve:

(a) This solution contains barium ions and acetate ions. The cation, Ba^{2+} , is an ion of one of the heavy alkaline earth metals and will therefore not affect the pH (summary point 4). The anion, CH_3COO^- , is the conjugate base of the weak acid CH_3COOH and will hydrolyze to produce OH^- ions, thereby making the solution basic (summary point 2).

(b) This solution contains NH_4^+ and Cl^- ions. NH_4^+ is the conjugate acid of a weak base (NH_3) and is therefore acidic (summary point 3). Cl^- is the conjugate base of a strong acid (HCl) and therefore has no influence on the pH of the solution (summary point 1). Because the solution contains an ion that is acidic (NH_4^+) and one that has no influence on pH (Cl^-), the solution of NH_4Cl will be acidic.

(c) This solution contains CH_3NH_3^+ and Br^- ions. CH_3NH_3^+ is the conjugate acid of a weak base (CH_3NH_2 , an amine) and is therefore acidic (summary point 3). Br^- is the conjugate base of a strong acid (HBr) and is therefore pH-neutral (summary point 1). Because the solution contains one ion that is acidic and one that is neutral, the solution of $\text{CH}_3\text{NH}_3\text{Br}$ will be acidic.

(d) This solution contains the K^+ ion, which is a cation of group 1A, and the NO_3^- ion, which is the conjugate base of the strong acid HNO_3 . Neither of the ions will react with water to any appreciable extent (summary points 1 and 4), making the solution neutral.

(e) This solution contains Al^{3+} and ClO_4^- ions. Cations, such as Al^{3+} , that are not in groups 1A or 2A are acidic (summary point 5). The ClO_4^- ion is the conjugate base of a strong acid (HClO_4) and therefore does not affect pH (summary point 1). Thus, the solution of $\text{Al}(\text{ClO}_4)_3$ will be acidic.

PRACTICE EXERCISE

In each of the following, indicate which salt in each of the following pairs will form the more acidic (or less basic) 0.010 M solution: (a) NaNO_3 , or $\text{Fe}(\text{NO}_3)_3$; (b) KBr , or KBrO ; (c) $\text{CH}_3\text{NH}_3\text{Cl}$, or BaCl_2 ; (d) NH_4NO_2 , or NH_4NO_3 .

Answers: (a) $\text{Fe}(\text{NO}_3)_3$, (b) KBr , (c) $\text{CH}_3\text{NH}_3\text{Cl}$, (d) NH_4NO_3



SAMPLE EXERCISE 16.19 | Predicting Whether the Solution of an Amphiprotic Anion Is Acidic or Basic

Predict whether the salt Na_2HPO_4 will form an acidic solution or a basic solution on dissolving in water.

SOLUTION

Analyze: We are asked to predict whether a solution of Na_2HPO_4 will be acidic or basic. This substance is an ionic compound composed of Na^+ and HPO_4^{2-} ions.

Plan: We need to evaluate each ion, predicting whether each is acidic or basic. Because Na^+ is a cation of group 1A, we know that it has no influence on pH. It is merely a spectator ion in acid–base chemistry. Thus, our analysis of whether the solution is acidic or basic must focus on the behavior of the HPO_4^{2-} ion. We need to consider the fact that HPO_4^{2-} can act as either an acid or a base.



The reaction with the larger equilibrium constant will determine whether the solution is acidic or basic.

Solve: The value of K_a for Equation 16.45, as shown in Table 16.3, is 4.2×10^{-13} . We must calculate the value of K_b for Equation 16.46 from the value of K_a for its conjugate acid, H_2PO_4^- . We make use of the relationship shown in Equation 16.40.

$$K_a \times K_b = K_w$$

We want to know K_b for the base HPO_4^{2-} , knowing the value of K_a for the conjugate acid H_2PO_4^- :

$$K_b(\text{HPO}_4^{2-}) \times K_a(\text{H}_2\text{PO}_4^-) = K_w = 1.0 \times 10^{-14}$$

Because K_a for H_2PO_4^- is 6.2×10^{-8} (Table 16.3), we calculate K_b for HPO_4^{2-} to be 1.6×10^{-7} . This is more than 10^5 times larger than K_a for HPO_4^{2-} ; thus, the reaction shown in Equation 16.46 predominates over that in Equation 16.45, and the solution will be basic.

■ PRACTICE EXERCISE

Predict whether the dipotassium salt of citric acid ($\text{K}_2\text{HC}_6\text{H}_5\text{O}_7$) will form an acidic or basic solution in water (see Table 16.3 for data).

Answer: acidic

16.10 ACID–BASE BEHAVIOR AND CHEMICAL STRUCTURE

When a substance is dissolved in water, it may behave as an acid, behave as a base, or exhibit no acid–base properties. How does the chemical structure of a substance determine which of these behaviors is exhibited by the substance? For example, why do some substances that contain OH groups behave as bases, releasing OH^- ions into solution, whereas others behave as acids, ionizing to release H^+ ions? Why are some acids stronger than others? In this section we will discuss briefly the effects of chemical structure on acid–base behavior.

Factors That Affect Acid Strength

A molecule containing H will transfer a proton only if the H—X bond is polarized in the following way:



In ionic hydrides, such as NaH, the reverse is true; the H atom possesses a negative charge and behaves as a proton acceptor. Essentially nonpolar H—X bonds, such as the H—C bond in CH_4 , produce neither acidic nor basic aqueous solutions.

A second factor that helps determine whether a molecule containing an H—X bond will donate a proton is the strength of the bond. Very strong bonds are less easily dissociated than weaker ones. This factor is important, for example, in the case of the hydrogen halides. The H—F bond is the most polar H—X bond. You therefore might expect that HF would be a very strong acid if the first factor were all that mattered. However, HF has the highest bond strength among the hydrogen halides, as seen in Table 8.4. As a result, HF is a weak acid, whereas all the other hydrogen halides are strong acids in water.

A third factor that affects the ease with which a hydrogen atom ionizes from HX is the stability of the conjugate base, X^- . In general, the greater the stability of the conjugate base, the stronger is the acid. The strength of an acid is often a combination of all three factors: (1) the polarity of the H—X bond, (2) the strength of the H—X bond, and (3) the stability of the conjugate base, X^- .

Binary Acids

In general, the H—X bond strength is the most important factor determining acid strength among the binary acids (those containing hydrogen and just one other element) in which X is in the same *group* in the periodic table. The strength of an H—X bond tends to decrease as the element X increases in size. As a result, the bond strength decreases and the acidity increases down a group. Thus, HCl is a stronger acid than HF, and H_2S is a stronger acid than H_2O .

	GROUP			
	4A	5A	6A	7A
Period 2	CH ₄ No acid or base properties	NH ₃ Weak base	H ₂ O ---	HF Weak acid
Period 3	SiH ₄ No acid or base properties	PH ₃ Weak base	H ₂ S Weak acid	HCl Strong acid

▲ **Figure 16.12 Trends in acid-base properties of binary hydrides.** The acidity of the binary compounds of hydrogen and nonmetals increases moving left to right across a period and moving top to bottom down a group.


Bond strengths change less moving across a row in the periodic table than they do down a group. As a result, bond polarity is the major factor determining acidity for binary acids in the same *row*. Thus, acidity increases as the electronegativity of the element X increases, as it generally does moving from left to right in a row. For example, the acidity of the second-row elements varies in the following order: CH₄ < NH₃ ≪ H₂O < HF. Because the C—H bond is essentially nonpolar, CH₄ shows no tendency to form H⁺ and CH₃[−] ions. Although the N—H bond is polar, NH₃ has a nonbonding pair of electrons on the nitrogen atom that dominates its chemistry, so NH₃ acts as a base rather than as an acid. The periodic trends in the acid strengths of binary compounds of hydrogen and the nonmetals of periods 2 and 3 are summarized in Figure 16.12 ▲.

GIVE IT SOME THOUGHT

What is the major factor determining the increase in acidity of binary acids going down a column of the periodic table? What is the major factor going across a period?


Oxyacids

Many common acids, such as sulfuric acid, contain one or more O—H bonds:




Acids in which OH groups and possibly additional oxygen atoms are bound to a central atom are called **oxyacids**. The OH group is also present in bases. What factors determine whether an OH group will behave as a base or as an acid?

Let's consider an OH group bound to some atom Y, which might in turn have other groups attached to it:



At one extreme, Y might be a metal, such as Na, K, or Mg. Because of their low electronegativities, the pair of electrons shared between Y and O is completely transferred to oxygen, and an ionic compound containing OH[−] is formed. Such compounds are therefore sources of OH[−] ions and behave as bases.




◀ **Figure 16.13 The acidity of oxyacids increases with increasing electronegativity of the central atom.** As the electronegativity of the atom attached to an OH group increases, the ease with which the hydrogen ion is released increases. The drift of electron density toward the electronegative atom further polarizes the O—H bond, which favors ionization. In addition, the electronegative atom will help stabilize the conjugate base, which also leads to a stronger acid. Because Cl is more electronegative than I, HClO is a stronger acid than HIO.

When Y is a nonmetal, the bond to O is covalent and the substance does not readily lose OH^- . Instead, these compounds are either acidic or neutral. Generally, as the electronegativity of Y increases, so will the acidity of the substance. This happens for two reasons: First, as electron density is drawn toward Y, the O—H bond becomes weaker and more polar, thereby favoring loss of H^+ (Figure 16.13 ▲). Second, because the conjugate base is usually an anion, its stability generally increases as the electronegativity of Y increases.

Many oxyacids contain additional oxygen atoms bonded to the central atom Y. The additional electronegative oxygen atoms pull electron density from the O—H bond, further increasing its polarity. Increasing the number of oxygen atoms also helps stabilize the conjugate base by increasing its ability to “spread out” its negative charge. Thus, the strength of an acid will increase as additional electronegative atoms bond to the central atom Y.

We can summarize these ideas as two simple rules that relate the acid strength of oxyacids to the electronegativity of Y and to the number of groups attached to Y.

1. For oxyacids that have the same number of OH groups and the same number of O atoms, acid strength increases with increasing electronegativity of the central atom Y. For example, the strength of the hypohalous acids, which have the structure H—O—Y , increases as the electronegativity of Y increases (Table 16.6 ▶).
2. For oxyacids that have the same central atom Y, acid strength increases as the number of oxygen atoms attached to Y increases. For example, the strength of the oxyacids of chlorine steadily increases from hypochlorous acid (HClO) to perchloric acid (HClO_4):



Because the oxidation number of the central atom increases as the number of attached O atoms increases, this correlation can be stated in an equivalent way: In a series of oxyacids, the acidity increases as the oxidation number of the central atom increases.

TABLE 16.6 ■ Electronegativity Values (EN) of Y and Acid-Dissociation Constants

Acid	EN of Y	K_a
HClO	3.0	3.0×10^{-8}
HBrO	2.8	2.5×10^{-9}
HIO	2.5	2.3×10^{-11}

SAMPLE EXERCISE 16.20 Predicting Relative Acidities from Composition and Structure

Arrange the compounds in each of the following series in order of increasing acid strength: (a) AsH₃, HI, NaH, H₂O; (b) H₂SO₄, H₂SeO₃, H₂SeO₄.

SOLUTION

Analyze: We are asked to arrange two sets of compounds in order from weakest acid to strongest acid. In (a), the substances are binary compounds containing H, whereas in (b) the substances are oxyacids.

Plan: For the binary compounds in part (a), we will consider the electronegativities of As, I, Na, and O relative to H. A higher electronegativity will cause the H to have a higher partial positive charge, causing the compound to be more acidic. For the oxyacids in part (b), we will consider both the relative electronegativities of the central atom (S and Se) and the number of oxygen atoms bonded to the central atom.

Solve:

(a) Because Na is on the left side of the periodic table, we know that it has a very low electronegativity. As a result, the hydrogen in NaH carries a negative charge. Thus NaH should be the least acidic (most basic) compound on the list. Because arsenic is less electronegative than oxygen, we might expect that AsH₃ would be a weak base toward water. We would make the same prediction by an extension of the trends shown in Figure 16.12. Further, we expect that the binary hydrogen compounds of the halogens, as the most electronegative element in each period, will be acidic relative to water. In fact, HI is one of the strong acids in water. Thus the order of increasing acidity is NaH < AsH₃ < H₂O < HI.


(b) The acids H₂SO₄ and H₂SeO₄ have the same number of O atoms and OH groups. In such cases, the acid strength increases with increasing electronegativity of the central atom. Because S is more electronegative than Se, we predict that H₂SO₄ is more acidic than H₂SeO₄. Next, we can compare H₂SeO₄ and H₂SeO₃. For acids with the same central atom, the acidity increases as the number of oxygen atoms bonded to the central atom increases. Thus, H₂SeO₄ should be a stronger acid than H₂SeO₃. Thus, we predict the order of increasing acidity to be H₂SeO₃ < H₂SeO₄ < H₂SO₄.

PRACTICE EXERCISE


In each of the following pairs choose the compound that leads to the more acidic (or less basic) solution: (a) HBr, HF; (b) PH₃, H₂S; (c) HNO₂, HNO₃; (d) H₂SO₃, H₂SeO₃.
Answers: (a) HBr; (b) H₂S; (c) HNO₃; (d) H₂SO₃

Carboxylic Acids

Another large group of acids is illustrated by acetic acid:




The portion of the structure shown in blue is called the *carboxyl group*, which is often written as COOH. Thus, the chemical formula of acetic acid is written as CH₃COOH, where only the hydrogen atom in the carboxyl group can be ionized. Acids that contain a carboxyl group are called **carboxylic acids**, and they form the largest category of organic acids. Formic acid and benzoic acid, whose structures are drawn below, are further examples of this large and important category of acids.




Acetic acid (CH₃COOH) is a weak acid ($K_a = 1.8 \times 10^{-5}$). Two factors contribute to the acidic behavior of carboxylic acids. First, the additional oxygen atom attached to the carbon of the carboxyl group draws electron density from the O—H bond, increasing its polarity and helping to stabilize the conjugate base.



Amino acids are the building blocks of proteins. The general structure of amino acids is shown here, where different amino acids have different R groups attached to the central carbon atom:



For example, in *glycine*, which is the simplest amino acid, R is a hydrogen atom, whereas in *alanine*, R is a CH₃ group.




Amino acids contain a carboxyl group and can therefore serve as acids. They also contain an NH₂ group, characteristic of amines (Section 16.7), and thus they can also act as bases. Amino acids, therefore, are amphoteric. For glycine, we might expect that the acid and the base reactions with water would be as follows:



The pH of a solution of glycine in water is about 6.0, indicating that it is a slightly stronger acid than a base.

The acid–base chemistry of amino acids is somewhat more complicated than shown in Equations 16.47 and 16.48, however. Because the COOH can act as an acid and the NH₂ group can act as a base, amino acids undergo a “self-contained” Brønsted–Lowry acid–base reaction in which the proton of the carboxyl group is transferred to the basic nitrogen atom:



Although the form of the amino acid on the right side of Equation 16.49 is electrically neutral overall, it has a positively charged end and a negatively charged end. A molecule of this type is called a *zwitterion* (German for “hybrid ion”).


Do amino acids exhibit any properties indicating that they behave as zwitterions? If so, they should behave similar to ionic substances. ∞ (Section 8.2) Crystalline amino acids (Figure 16.14 ▼) have relatively high melting points, usually above 200 °C, which is characteristic of ionic solids. Amino acids are far more soluble in water than in nonpolar solvents. In addition, the dipole moments of amino acids are large, consistent with a large separation of charge in the molecule. Thus, the ability of amino acids to act simultaneously as acids and bases has important effects on their properties.

Related Exercise: 16.119



▲ **Figure 16.14 Lysine.** One of the amino acids found in proteins, lysine is available as a dietary supplement. The L on the label refers to a specific arrangement of atoms that is found in naturally occurring amino acids. Molecules with the L arrangement are mirror images of molecules with the D arrangement, much like our left hand is a mirror image of our right hand.

Second, the conjugate base of a carboxylic acid (a *carboxylate anion*) can exhibit resonance (Section 8.6), which contributes further to the stability of the anion by spreading the negative charge over several atoms:




GIVE IT SOME THOUGHT

What group of atoms is present in all carboxylic acids?


16.11 LEWIS ACIDS AND BASES

For a substance to be a proton acceptor (a Brønsted–Lowry base), it must have an unshared pair of electrons for binding the proton. NH_3 , for example, acts as a proton acceptor. Using Lewis structures, we can write the reaction between H^+ and NH_3 as follows:



G. N. Lewis was the first to notice this aspect of acid–base reactions. He proposed a definition of acid and base that emphasizes the shared electron pair: A **Lewis acid** is an electron-pair acceptor, and a **Lewis base** is an electron-pair donor.

Every base that we have discussed thus far—whether it be OH^- , H_2O , an amine, or an anion—is an electron-pair donor. Everything that is a base in the Brønsted–Lowry sense (a proton acceptor) is also a base in the Lewis sense (an electron-pair donor). In the Lewis theory, however, a base can donate its electron pair to something other than H^+ . The Lewis definition therefore greatly increases the number of species that can be considered acids; H^+ is a Lewis acid, but not the only one. For example, consider the reaction between NH_3 and BF_3 . This reaction occurs because BF_3 has a vacant orbital in its valence shell. ∞ (Section 8.7) It therefore acts as an electron-pair acceptor (a Lewis acid) toward NH_3 , which donates the electron pair. The curved arrow shows the donation of a pair of electrons from N to B to form a covalent bond:




GIVE IT SOME THOUGHT

What feature must any molecule or ion have to act as a Lewis base?


Our emphasis throughout this chapter has been on water as the solvent and on the proton as the source of acidic properties. In such cases we find the Brønsted–Lowry definition of acids and bases to be the most useful. In fact, when we speak of a substance as being acidic or basic, we are usually thinking of aqueous solutions and using these terms in the Arrhenius or Brønsted–Lowry sense. The advantage of the Lewis theory is that it allows us to treat a wider variety of reactions, including those that do not involve proton transfer, as acid–base reactions. To avoid confusion, a substance such as BF_3 is rarely called an acid unless it is clear from the context that we are using the term in the sense of the Lewis definition. Instead, substances that function as electron-pair acceptors are referred to explicitly as “Lewis acids.”

Lewis acids include molecules that, like BF_3 , have an incomplete octet of electrons. In addition, many simple cations can function as Lewis acids. For example, Fe^{3+} interacts strongly with cyanide ions to form the ferricyanide ion, $[\text{Fe}(\text{CN})_6]^{3-}$.



The Fe^{3+} ion has vacant orbitals that accept the electron pairs donated by the cyanide ions; we will learn more in Chapter 24 about just which orbitals are used by the Fe^{3+} ion. The metal ion is highly charged, too, which contributes to the interaction with CN^- ions.

Some compounds with multiple bonds can behave as Lewis acids. For example, the reaction of carbon dioxide with water to form carbonic acid (H_2CO_3) can be pictured as an attack by a water molecule on CO_2 , in which the water acts as an electron-pair donor and the CO_2 as an electron-pair acceptor, as shown in the margin. The electron pair of one of the carbon–oxygen double bonds is moved onto the oxygen, leaving a vacant orbital on the carbon that can act as an electron-pair acceptor. We have shown the shift of these electrons with arrows. After forming the initial acid–base product, a proton moves from one oxygen to another, thereby forming carbonic acid. A similar kind of Lewis acid–base reaction takes place when any oxide of a nonmetal dissolves in water to form an acidic solution.



Hydrolysis of Metal Ions


As we have already seen, most metal ions behave as acids in aqueous solution. ∞ (Section 16.9) For example, an aqueous solution of $\text{Fe}(\text{NO}_3)_3$ is quite acidic. An aqueous solution of ZnCl_2 is also acidic, though to a lesser extent. The Lewis concept helps explain the interactions between metal ions and water molecules that give rise to this acidic behavior.

Because metal ions are positively charged, they attract the unshared electron pairs of water molecules. It is primarily this interaction, referred to as *hydration*, that causes salts to dissolve in water. ∞ (Section 13.1) The process of hydration can be thought of as a Lewis acid–base interaction in which the metal ion acts as a Lewis acid and the water molecules as Lewis bases. When a water molecule interacts with the positively charged metal ion, electron density is drawn from the oxygen, as illustrated in Figure 16.15. This flow of electron density causes the O—H bond to become more polarized; as a result, water molecules bound to the metal ion are more acidic than those in the bulk solvent.

The hydrated Fe^{3+} ion, $\text{Fe}(\text{H}_2\text{O})_6^{3+}$, which we usually represent simply as $\text{Fe}^{3+}(\text{aq})$, acts as a source of protons:



The acid-dissociation constant for this hydrolysis reaction has the value $K_a = 2 \times 10^{-3}$, so $\text{Fe}^{3+}(\text{aq})$ is a fairly strong acid. Acid-dissociation constants for hydrolysis reactions generally increase with increasing charge and decreasing radius of the ion (Figure 16.15). Thus, the Cu^{2+} ion, which has a smaller charge and a larger radius than Fe^{3+} , forms less acidic solutions than Fe^{3+} . The K_a for $\text{Cu}^{2+}(\text{aq})$ is 1×10^{-8} . The acid hydrolysis of a number of salts of metal ions is demonstrated in Figure 16.16. Note that the Na^+ ion, which is large and has only a 1+ charge (and which we have previously identified as the cation of a strong base), exhibits no acid hydrolysis and yields a neutral solution.



▲ **Figure 16.15 The acidity of a hydrated cation depends on cation charge and size.** The interaction between a water molecule and a cation is much stronger when the cation is a smaller ion of higher charge. The pull of electron density toward the cation weakens the polar O—H bond of the water molecule and allows the transfer of a H^+ ion to a nearby water molecule. As a result, hydrated cations tend to be acidic, with their acidities increasing with increasing charge and decreasing size.

Salt:	NaNO_3	$\text{Ca}(\text{NO}_3)_2$	$\text{Zn}(\text{NO}_3)_2$	$\text{Al}(\text{NO}_3)_3$
Indicator:	Bromthymol blue	Bromthymol blue	Methyl red	Methyl orange
Estimated pH:	7.0	6.9	5.5	3.5


◀ **Figure 16.16 Effect of cations on the pH of a solution.** The pH values of 1.0 M solutions of a series of nitrate salts are estimated using acid–base indicators.

GIVE IT SOME THOUGHT

Which of the following cations will be most acidic and why: Ca^{2+} , Fe^{2+} , Fe^{3+} ?

SAMPLE INTEGRATIVE EXERCISE | Putting Concepts Together

Phosphorous acid (H_3PO_3) has the following Lewis structure.



(a) Explain why H_3PO_3 is diprotic and not triprotic. (b) A 25.0-mL sample of a solution of H_3PO_3 is titrated with 0.102 M NaOH. It requires 23.3 mL of NaOH to neutralize both acidic protons. What is the molarity of the H_3PO_3 solution? (c) The original solution from part (b) has a pH of 1.59. Calculate the percent ionization and K_{a1} for H_3PO_3 , assuming that $K_{a1} \gg K_{a2}$. (d) How does the osmotic pressure of a 0.050 M solution of HCl compare qualitatively with that of a 0.050 M solution of H_3PO_3 ? Explain.

SOLUTION

We will use what we have learned about molecular structure and its impact on acidic behavior to answer part (a). We will then use stoichiometry and the relationship between pH and $[\text{H}^+]$ to answer parts (b) and (c). Finally, we will consider acid strength in order to compare the colligative properties of the two solutions in part (d).

(a) Acids have polar H—X bonds. From Figure 8.6 we see that the electronegativity of H is 2.1 and that of P is also 2.1. Because the two elements have the same electronegativity, the H—P bond is nonpolar. ∞ (Section 8.4) Thus, this H cannot be acidic. The other two H atoms, however, are bonded to O, which has an electronegativity of 3.5. The H—O bonds are therefore polar, with H having a partial positive charge. These two H atoms are consequently acidic.

(b) The chemical equation for the neutralization reaction is



From the definition of molarity, $M = \text{mol/L}$, we see that moles = $M \times L$. ∞ (Section 4.5) Thus, the number of moles of NaOH added to the solution is $(0.0233 \text{ L})(0.102 \text{ mol/L}) = 2.38 \times 10^{-3} \text{ mol NaOH}$. The balanced equation indicates that 2 mol of NaOH is consumed for each mole of H_3PO_3 . Thus, the number of moles of H_3PO_3 in the sample is


$$(2.38 \times 10^{-3} \text{ mol NaOH}) \left(\frac{1 \text{ mol H}_3\text{PO}_3}{2 \text{ mol NaOH}} \right) = 1.19 \times 10^{-3} \text{ mol H}_3\text{PO}_3$$

The concentration of the H_3PO_3 solution, therefore, equals $(1.19 \times 10^{-3} \text{ mol}) / (0.0250 \text{ L}) = 0.0476 \text{ M}$.

(c) From the pH of the solution, 1.59, we can calculate $[\text{H}^+]$ at equilibrium.

$$[\text{H}^+] = \text{antilog}(-1.59) = 10^{-1.59} = 0.026 \text{ M (two significant figures)}$$

Because $K_{a1} \gg K_{a2}$, the vast majority of the ions in solution are from the first ionization step of the acid.



Because one H_2PO_3^- ion forms for each H^+ ion formed, the equilibrium concentrations of H^+ and H_2PO_3^- are equal: $[\text{H}^+] = [\text{H}_2\text{PO}_3^-] = 0.026 \text{ M}$. The equilibrium concentration of H_3PO_3 equals the initial concentration minus the amount that ionizes to form H^+ and H_2PO_3^- : $[\text{H}_3\text{PO}_3] = 0.0476 \text{ M} - 0.026 \text{ M} = 0.022 \text{ M}$ (two significant figures). These results can be tabulated as follows:

	$\text{H}_3\text{PO}_3(aq)$	\rightleftharpoons	$\text{H}^+(aq)$	+	$\text{H}_2\text{PO}_3^-(aq)$
Initial	0.0476 M		0		0
Change	-0.026 M		+0.026 M		+0.026 M
Equilibrium	0.022 M		0.026 M		0.026 M

The percent ionization is

$$\text{Percent ionization} = \frac{[\text{H}^+]_{\text{equilibrium}}}{[\text{H}_3\text{PO}_3]_{\text{initial}}} \times 100\% = \frac{0.026 \text{ M}}{0.0476 \text{ M}} \times 100\% = 55\%$$

The first acid-dissociation constant is

$$K_{a1} = \frac{[\text{H}^+][\text{H}_2\text{PO}_3^-]}{[\text{H}_3\text{PO}_3]} = \frac{(0.026)(0.026)}{0.022} = 0.031$$

(d) Osmotic pressure is a colligative property and depends on the total concentration of particles in solution. ∞ (Section 13.5) Because HCl is a strong acid, a 0.050 M solution will contain 0.050 M $\text{H}^+(aq)$ and 0.050 M $\text{Cl}^-(aq)$, or a total of 0.100 mol/L of particles. Because H_3PO_3 is a weak acid, it ionizes to a lesser extent than HCl, and, hence, there are fewer particles in the H_3PO_3 solution. As a result, the H_3PO_3 solution will have the lower osmotic pressure.

CHAPTER REVIEW

SUMMARY AND KEY TERMS

Section 16.1 Acids and bases were first recognized by the properties of their aqueous solutions. For example, acids turn litmus red, whereas bases turn litmus blue. Arrhenius recognized that the properties of acidic solutions are due to $\text{H}^+(aq)$ ions and those of basic solutions are due to $\text{OH}^-(aq)$ ions.

Section 16.2 The Brønsted–Lowry concept of acids and bases is more general than the Arrhenius concept and emphasizes the transfer of a proton (H^+) from an acid to a base. The H^+ ion, which is merely a proton with no surrounding valence electrons, is strongly bound to water. For this reason, the **hydronium ion**, $\text{H}_3\text{O}^+(aq)$, is often used to represent the predominant form of H^+ in water instead of the simpler $\text{H}^+(aq)$.

A **Brønsted–Lowry acid** is a substance that donates a proton to another substance; a **Brønsted–Lowry base** is a substance that accepts a proton from another substance. Water is an **amphiprotic** substance, one that can function as either a Brønsted–Lowry acid or base, depending on the substance with which it reacts.

The **conjugate base** of a Brønsted–Lowry acid is the species that remains when a proton is removed from the acid. The **conjugate acid** of a Brønsted–Lowry base is the species formed by adding a proton to the base. Together, an acid and its conjugate base (or a base and its conjugate acid) are called a **conjugate acid–base pair**.

The acid–base strengths of conjugate acid–base pairs are related: The stronger an acid, the weaker is its conjugate base; the weaker an acid, the stronger is its conjugate base. In every acid–base reaction, the position of the equilibrium favors the transfer of the proton from the stronger acid to the stronger base.

Section 16.3 Water ionizes to a slight degree, forming $\text{H}^+(aq)$ and $\text{OH}^-(aq)$. The extent of this **autoionization** is expressed by the **ion-product constant** for water:

$K_w = [\text{H}^+][\text{OH}^-] = 1.0 \times 10^{-14}$ (25 °C). This relationship describes both pure water and aqueous solutions. The K_w expression indicates that the product of $[\text{H}^+]$ and $[\text{OH}^-]$ is a constant. Thus, as $[\text{H}^+]$ increases, $[\text{OH}^-]$ decreases. Acidic solutions are those that contain more $\text{H}^+(aq)$ than $\text{OH}^-(aq)$; basic solutions contain more $\text{OH}^-(aq)$ than $\text{H}^+(aq)$.

Section 16.4 The concentration of $\text{H}^+(aq)$ can be expressed in terms of **pH**: $\text{pH} = -\log[\text{H}^+]$. At 25 °C the pH of a neutral solution is 7.00, whereas the pH of an acidic solution is below 7.00, and the pH of a basic solution is above 7.00. The pX notation is also used to represent the negative log of other small quantities, as in pOH and $\text{p}K_w$. The pH of a solution can be measured using a pH meter, or it can be estimated using acid–base indicators.

Section 16.5 Strong acids are strong electrolytes, ionizing completely in aqueous solution. The common strong acids are HCl, HBr, HI, HNO_3 , HClO_3 , HClO_4 , and H_2SO_4 . The conjugate bases of strong acids have negligible basicity.

Common strong bases are the ionic hydroxides of alkali metals and the heavy alkaline earth metals. The cations of these metals have negligible acidity.

Section 16.6 Weak acids are weak electrolytes; only some of the molecules exist in solution in ionized form. The extent of ionization is expressed by the **acid-dissociation constant**, K_a , which is the equilibrium constant for the reaction $\text{HA}(aq) \rightleftharpoons \text{H}^+(aq) + \text{A}^-(aq)$, which can also be written $\text{HA}(aq) + \text{H}_2\text{O}(l) \rightleftharpoons \text{H}_3\text{O}^+(aq) + \text{A}^-(aq)$. The larger the value of K_a , the stronger is the acid. For solutions of the same concentration, a stronger acid also has a larger **percent ionization**. The concentration of a weak acid and its K_a value can be used to calculate the pH of a solution.

Polyprotic acids, such as H_2SO_3 , have more than one ionizable proton. These acids have acid-dissociation constants that decrease in magnitude in the order

$K_{a1} > K_{a2} > K_{a3}$. Because nearly all the $H^+(aq)$ in a polyprotic acid solution comes from the first dissociation step, the pH can usually be estimated satisfactorily by considering only K_{a1} .

Sections 16.7 and 16.8 Weak bases include NH_3 , amines, and the anions of weak acids. The extent to which a weak base reacts with water to generate the corresponding conjugate acid and OH^- is measured by the **base-dissociation constant**, K_b . This is the equilibrium constant for the reaction $B(aq) + H_2O(l) \rightleftharpoons HB^+(aq) + OH^-(aq)$, where B is the base.

The relationship between the strength of an acid and the strength of its conjugate base is expressed quantitatively by the equation $K_a \times K_b = K_w$, where K_a and K_b are dissociation constants for conjugate acid–base pairs.

Section 16.9 The acid–base properties of salts can be ascribed to the behavior of their respective cations and anions. The reaction of ions with water, with a resultant change in pH, is called **hydrolysis**. The cations of the alkali metals and the alkaline earth metals and the anions of strong acids do not undergo hydrolysis. They are always spectator ions in acid–base chemistry.

Section 16.10 The tendency of a substance to show acidic or basic characteristics in water can be correlated with its chemical structure. Acid character requires the presence of a highly polar $H-X$ bond. Acidity is also favored when the $H-X$ bond is weak and when the X^- ion is very stable.

For **oxyacids** with the same number of OH groups and the same number of O atoms, acid strength increases with increasing electronegativity of the central atom. For oxyacids with the same central atom, acid strength increases as the number of oxygen atoms attached to the central atom increases. The structures of **carboxylic acids**, which are organic acids containing the COOH group, also help us to understand their acidity.

Section 16.11 The Lewis concept of acids and bases emphasizes the shared electron pair rather than the proton. A **Lewis acid** is an electron-pair acceptor, and a **Lewis base** is an electron-pair donor. The Lewis concept is more general than the Brønsted–Lowry concept because it can apply to cases in which the acid is some substance other than H^+ . The Lewis concept helps to explain why many hydrated metal cations form acidic aqueous solutions. The acidity of these cations generally increases as their charge increases and as the size of the metal ion decreases.

KEY SKILLS

- Understand the nature of the hydrated proton, represented as either $H^+(aq)$ or $H_3O^+(aq)$.
- Define and identify Arrhenius acids and bases.
- Define and identify Brønsted–Lowry acids and bases, and identify conjugate acid–base pairs.
- Relate the strength of an acid to the strength of its conjugate base.
- Understand how the equilibrium position of a proton transfer reaction relates the strengths of the acids and bases involved.
- Describe the autoionization of water and understand how $[H_3O^+]$ and $[OH^-]$ are related.
- Calculate the pH of a solution given $[H_3O^+]$ or $[OH^-]$.
- Calculate the pH of a strong acid or strong base given its concentration.
- Calculate K_a or K_b for a weak acid or weak base given its concentration and the pH of the solution.
- Calculate the pH of a weak acid or weak base or its percent ionization given its concentration and K_a or K_b .
- Calculate K_b for a weak base given K_a of its conjugate acid, and similarly calculate K_a from K_b .
- Predict whether an aqueous solution of a salt will be acidic, basic, or neutral.
- Predict the relative strength of a series of acids from their molecular structures.
- Define and identify Lewis acids and bases.


KEY EQUATIONS

- $K_w = [H_3O^+][OH^-] = [H^+][OH^-] = 1.0 \times 10^{-14}$ [16.16] The ion product of water at 25 °C
- $pH = -\log[H^+]$ [16.17] Definition of pH
- $pOH = -\log[OH^-]$ [16.18] Definition of pOH

- $\text{pH} + \text{pOH} = 14.00$ [16.20] Relationship between pH and pOH
- $K_a = \frac{[\text{H}_3\text{O}^+][\text{A}^-]}{[\text{HA}]}$ or $K_a = \frac{[\text{H}^+][\text{A}^-]}{[\text{HA}]}$ [16.25] The acid dissociation constant for a weak acid, HA
- Percent ionization = $\frac{[\text{H}^+]_{\text{equilibrium}}}{[\text{HA}]_{\text{initial}}} \times 100\%$ [16.27] Percent ionization of a weak acid
- $K_b = \frac{[\text{BH}^+][\text{OH}^-]}{[\text{B}]}$ [16.34] The base dissociation constant for a weak base, B
- $K_a \times K_b = K_w$ [16.40] The relationship between the acid and base dissociation constants of a conjugate acid–base pair


VISUALIZING CONCEPTS

- 16.1 (a) Identify the Brønsted–Lowry acid and the Brønsted–Lowry base in the following reaction:




- (b) Identify the Lewis acid and the Lewis base in the reaction. [Sections 16.2 and 16.11]

- 16.2 The following diagrams represent aqueous solutions of two monoprotic acids, HA (A = X or Y). The water molecules have been omitted for clarity. (a) Which is the stronger acid, HX or HY? (b) Which is the stronger base, X^- or Y^- ? (c) If you mix equal concentrations of HX and NaY, will the equilibrium




- lie mostly to the right ($K_c > 1$) or to the left ($K_c < 1$)? [Section 16.2]




- 16.3 The following diagrams represent aqueous solutions of three acids, HX, HY, and HZ. The water molecules have been omitted for clarity, and the hydrated proton is represented as a simple sphere rather than as a hydronium ion. (a) Which of the acids is a strong acid? Explain.

- (b) Which acid would have the smallest acid-dissociation constant, K_a ? (c) Which solution would have the highest pH? [Sections 16.5 and 16.6]



- 16.4 In which of the following cases is the approximation that the equilibrium concentration of $\text{H}^+(aq)$ is small relative to the initial concentration of HA likely to be most valid: (a) initial $[\text{HA}] = 0.100 \text{ M}$ and $K_a = 1.0 \times 10^{-6}$, (b) initial $[\text{HA}] = 0.100 \text{ M}$ and $K_a = 1.0 \times 10^{-4}$, (c) initial $[\text{HA}] = 0.100 \text{ M}$ and $K_a = 1.0 \times 10^{-3}$? [Section 16.6]

- 16.5 (a) Which of these three lines represents the effect of concentration on the percent ionization of a weak acid? (b) Explain in qualitative terms why the curve you choose has the shape it does. [Section 16.6]




- 16.3 The following diagrams represent aqueous solutions of three acids, HX, HY, and HZ. The water molecules have been omitted for clarity, and the hydrated proton is represented as a simple sphere rather than as a hydronium ion. (a) Which of the acids is a strong acid? Explain.

16.6 Refer to the diagrams accompanying Exercise 16.3. (a) Rank the anions, X^- , Y^- , and Z^- , in order of increasing basicity. (b) Which of the ions would have the largest base-dissociation constant, K_b ? [Sections 16.2 and 16.8]


16.7 (a) Draw the Lewis structure for the following molecule and explain why it is able to act as a base. (b) To what class of organic compounds does this substance belong? (See the color key in Exercise 16.1.) [Section 16.7]




16.8 The following diagram represents an aqueous solution formed by dissolving a sodium salt of a weak acid in water. The diagram shows only the Na^+ ions, the X^- ions, and the HX molecules. What ion is missing from the diagram? If the drawing is completed by drawing all the ions, how many of the missing ions should be shown? [Section 16.9]



16.9 (a) What kinds of acids are represented by the following molecular models? (b) Indicate how the acidity of each molecule is affected by increasing the electronegativity of the atom X, and explain the origin of the effect. [Section 16.10]




16.10 In this model of acetylsalicylic acid (aspirin), identify the carboxyl group in the molecule. [Section 16.10]




16.11 Rank the following acids in order of increasing acidity: CH_3COOH , CH_2ClCOOH , CHCl_2COOH , CCl_3COOH , CF_3COOH . [Section 16.10]

16.12 (a) The following diagram represents the reaction of PCl_4^+ with Cl^- . Draw the Lewis structures for the reactants and products, and identify the Lewis acid and the Lewis base in the reaction.



(b) The following reaction represents the acidity of a hydrated cation. How does the equilibrium constant for the reaction change as the charge of the cation increases? [Section 16.11]



EXERCISES

Arrhenius and Brønsted–Lowry Acids and Bases

16.13 Although HCl and H_2SO_4 have very different properties as pure substances, their aqueous solutions possess many common properties. List some general properties of these solutions, and explain their common behavior in terms of the species present.

16.14 Although pure NaOH and NH_3 have very different properties, their aqueous solutions possess many common properties. List some general properties of these solutions, and explain their common behavior in terms of the species present.

16.15 (a) What is the difference between the Arrhenius and the Brønsted–Lowry definitions of an acid? (b) $\text{NH}_3(\text{g})$ and $\text{HCl}(\text{g})$ react to form the ionic solid $\text{NH}_4\text{Cl}(\text{s})$ (Figure 16.3). Which substance is the Brønsted–Lowry acid in this reaction? Which is the Brønsted–Lowry base?




16.16 (a) What is the difference between the Arrhenius and the Brønsted–Lowry definitions of a base? (b) When ammonia is dissolved in water, it behaves both as an Arrhenius base and as a Brønsted–Lowry base. Explain.

16.17 (a) Give the conjugate base of the following Brønsted–Lowry acids: (i) HIO_3 , (ii) NH_4^+ . (b) Give the conjugate acid of the following Brønsted–Lowry bases: (i) O^{2-} , (ii) H_2PO_4^- .




16.18 (a) Give the conjugate base of the following Brønsted–Lowry acids: (i) $\text{C}_6\text{H}_5\text{COOH}$. (ii) HPO_4^{2-} . (b) Give the conjugate acid of the following Brønsted–Lowry bases: (i) CO_3^{2-} , (ii) $\text{C}_2\text{H}_5\text{NH}_2$.

16.19 Designate the Brønsted–Lowry acid and the Brønsted–Lowry base on the left side of each of the following

equations, and also designate the conjugate acid and conjugate base on the right side:



16.20 Designate the Brønsted–Lowry acid and the Brønsted–Lowry base on the left side of each equation, and also designate the conjugate acid and conjugate base on the right side.



16.21 (a) The hydrogen oxalate ion (HC_2O_4^-) is amphiprotic. Write a balanced chemical equation showing how it acts as an acid toward water and another equation showing how it acts as a base toward water. (b) What is the conjugate acid of HC_2O_4^- ? What is its conjugate base?

16.22 (a) Write an equation for the reaction in which $\text{H}_2\text{C}_6\text{H}_7\text{O}_5^-(aq)$ acts as a base in $\text{H}_2\text{O}(l)$. (b) Write an equation for the reaction in which $\text{H}_2\text{C}_6\text{H}_7\text{O}_5^-(aq)$ acts as an acid in $\text{H}_2\text{O}(l)$. (c) What is the conjugate acid of $\text{H}_2\text{C}_6\text{H}_7\text{O}_5^-$? What is its conjugate base?

16.23 Label each of the following as being a strong base, a weak base, or a species with negligible basicity. In each case write the formula of its conjugate acid, and indicate whether the conjugate acid is a strong acid, a weak acid,




or a species with negligible acidity: (a) CH_3COO^- , (b) HCO_3^- , (c) O^{2-} , (d) Cl^- , (e) NH_3 .

16.24 Label each of the following as being a strong acid, a weak acid, or a species with negligible acidity. In each case write the formula of its conjugate base, and indicate whether the conjugate base is a strong base, a weak base, or a species with negligible basicity: (a) HNO_2 , (b) H_2SO_4 , (c) HPO_4^{2-} , (d) CH_4 , (e) CH_3NH_3^+ (an ion related to NH_4^+).




16.25 (a) Which of the following is the stronger Brønsted–Lowry acid, HBrO or HBr ? (b) Which is the stronger Brønsted–Lowry base, F^- or Cl^- ? Briefly explain your choices.

16.26 (a) Which of the following is the stronger Brønsted–Lowry acid, HNO_3 or HNO_2 ? (b) Which is the stronger Brønsted–Lowry base, NH_3 or H_2O ? Briefly explain your choices.

16.27 Predict the products of the following acid–base reactions, and predict whether the equilibrium lies to the left or to the right of the equation:



16.28 Predict the products of the following acid–base reactions, and predict whether the equilibrium lies to the left or to the right of the equation:



Autoionization of Water

16.29 (a) What does the term *autoionization* mean? (b) Explain why pure water is a poor conductor of electricity. (c) You are told that an aqueous solution is acidic. What does this statement mean?

16.30 (a) Write a chemical equation that illustrates the autoionization of water. (b) Write the expression for the ion-product constant for water, K_w . Why is $[\text{H}_2\text{O}]$ absent from this expression? (c) A solution is described as basic. What does this statement mean?

16.31 Calculate $[\text{H}^+]$ for each of the following solutions, and indicate whether the solution is acidic, basic, or neutral: (a) $[\text{OH}^-] = 0.00045 \text{ M}$; (b) $[\text{OH}^-] = 8.8 \times 10^{-9} \text{ M}$; (c) a solution in which $[\text{OH}^-]$ is 100 times greater than $[\text{H}^+]$.

16.32 Calculate $[\text{OH}^-]$ for each of the following solutions, and indicate whether the solution is acidic, basic, or neutral: (a) $[\text{H}^+] = 0.0045 \text{ M}$; (b) $[\text{H}^+] = 1.5 \times 10^{-9} \text{ M}$; (c) a solution in which $[\text{H}^+]$ is 10 times greater than $[\text{OH}^-]$.

16.33 At the freezing point of water (0°C), $K_w = 1.2 \times 10^{-15}$. Calculate $[\text{H}^+]$ and $[\text{OH}^-]$ for a neutral solution at this temperature.

16.34 Deuterium oxide (D_2O , where D is deuterium, the hydrogen-2 isotope) has an ion-product constant, K_w , of 8.9×10^{-16} at 20°C . Calculate $[\text{D}^+]$ and $[\text{OD}^-]$ for pure (neutral) D_2O at this temperature.

The pH Scale

16.35 By what factor does $[\text{H}^+]$ change for a pH change of (a) 2.00 units, (b) 0.50 units?

16.36 Consider two solutions, solution A and solution B. $[\text{H}^+]$ in solution A is 500 times greater than that in solution B. What is the difference in the pH values of the two solutions?

16.37 (a) If NaOH is added to water, how does $[\text{H}^+]$ change? How does pH change? (b) Use the pH values in Figure 16.5 to estimate the pH of a solution with $[\text{H}^+] = 0.0006 \text{ M}$. Is the solution acidic or basic? (c) If the pH of a solution is 5.2, first estimate and then calculate the molar concentrations of $\text{H}^+(aq)$ and $\text{OH}^-(aq)$ in the solution.

16.38 (a) If HNO_3 is added to water, how does $[\text{OH}^-]$ change? How does pH change? (b) Use the pH values in Figure 16.5 to estimate the pH of a solution with $[\text{OH}^-] = 0.014 \text{ M}$. Is the solution acidic or basic? (c) If $\text{pH} = 6.6$, first estimate and then calculate the molar concentrations of $\text{H}^+(\text{aq})$ and $\text{OH}^-(\text{aq})$ in the solution.

16.39 Complete the following table by calculating the missing entries and indicating whether the solution is acidic or basic.

$[\text{H}^+]$	$\text{OH}^-(\text{aq})$	pH	pOH	Acidic or basic?
$7.5 \times 10^{-3} \text{ M}$				
	$3.6 \times 10^{-10} \text{ M}$			
		8.25		
			5.70	

16.40 Complete the following table by calculating the missing entries. In each case indicate whether the solution is acidic or basic.

pH	pOH	$[\text{H}^+]$	$[\text{OH}^-]$	Acidic or basic?
11.25				
	6.02			
		$4.4 \times 10^{-4} \text{ M}$		
			$8.5 \times 10^{-3} \text{ M}$	

16.41 The average pH of normal arterial blood is 7.40. At normal body temperature (37°C), $K_w = 2.4 \times 10^{-14}$. Calculate $[\text{H}^+]$, $[\text{OH}^-]$, and pOH for blood at this temperature.

16.42 Carbon dioxide in the atmosphere dissolves in raindrops to produce carbonic acid (H_2CO_3), causing the pH of clean, unpolluted rain to range from about 5.2 to 5.6. What are the ranges of $[\text{H}^+]$ and $[\text{OH}^-]$ in the raindrops?

Strong Acids and Bases

16.43 (a) What is a strong acid? (b) A solution is labeled 0.500 M HCl . What is $[\text{H}^+]$ for the solution? (c) Which of the following are strong acids: HF , HCl , HBr , HI ?

16.44 (a) What is a strong base? (b) A solution is labeled $0.035 \text{ M Sr}(\text{OH})_2$. What is $[\text{OH}^-]$ for the solution? (c) Is the following statement true or false? Because $\text{Mg}(\text{OH})_2$ is not very soluble, it cannot be a strong base. Explain.

16.45 Calculate the pH of each of the following strong acid solutions: (a) $8.5 \times 10^{-3} \text{ M HBr}$, (b) 1.52 g of HNO_3 in 575 mL of solution, (c) 5.00 mL of 0.250 M HClO_4 diluted to 50.0 mL , (d) a solution formed by mixing 10.0 mL of 0.100 M HBr with 20.0 mL of 0.200 M HCl .

16.46 Calculate the pH of each of the following strong acid solutions: (a) 0.00135 M HNO_3 , (b) 0.425 g of HClO_4 in 2.00 L of solution, (c) 5.00 mL of 1.00 M HCl diluted to 0.500 L , (d) a mixture formed by adding 50.0 mL of 0.020 M HCl to 150 mL of 0.010 M HI .

16.47 Calculate $[\text{OH}^-]$ and pH for (a) $1.5 \times 10^{-3} \text{ M Sr}(\text{OH})_2$, (b) 2.250 g of LiOH in 250.0 mL of solution, (c) 1.00 mL of 0.175 M NaOH diluted to 2.00 L , (d) a solution formed by adding 5.00 mL of 0.105 M KOH to 15.0 mL of $9.5 \times 10^{-2} \text{ M Ca}(\text{OH})_2$.

16.48 Calculate $[\text{OH}^-]$ and pH for each of the following strong base solutions: (a) 0.082 M KOH , (b) 1.065 g of KOH in 500.0 mL of solution, (c) 10.0 mL of $0.0105 \text{ M Ca}(\text{OH})_2$ diluted to 500.0 mL , (d) a solution formed by mixing 10.0 mL of $0.015 \text{ M Ba}(\text{OH})_2$ with 40.0 mL of $7.5 \times 10^{-3} \text{ M NaOH}$.

16.49 Calculate the concentration of an aqueous solution of NaOH that has a pH of 11.50.

16.50 Calculate the concentration of an aqueous solution of $\text{Ca}(\text{OH})_2$ that has a pH of 12.05.

Weak Acids

16.51 Write the chemical equation and the K_a expression for the ionization of each of the following acids in aqueous solution. First show the reaction with $\text{H}^+(\text{aq})$ as a product and then with the hydronium ion: (a) HBrO_2 , (b) $\text{C}_2\text{H}_5\text{COOH}$.

16.52 Write the chemical equation and the K_a expression for the acid dissociation of each of the following acids in aqueous solution. First show the reaction with $\text{H}^+(\text{aq})$ as a product and then with the hydronium ion: (a) $\text{C}_6\text{H}_5\text{COOH}$, (b) HCO_3^- .

16.53 Lactic acid ($\text{CH}_3\text{CH}(\text{OH})\text{COOH}$) has one acidic hydrogen. A 0.10 M solution of lactic acid has a pH of 2.44. Calculate K_a .

16.54 Phenylacetic acid ($\text{C}_6\text{H}_5\text{CH}_2\text{COOH}$) is one of the substances that accumulates in the blood of people with phenylketonuria, an inherited disorder that can cause mental retardation or even death. A 0.085 M solution of $\text{C}_6\text{H}_5\text{CH}_2\text{COOH}$ has a pH of 2.68. Calculate the K_a value for this acid.

- 16.55** A 0.100 M solution of chloroacetic acid (ClCH_2COOH) is 11.0% ionized. Using this information, calculate $[\text{ClCH}_2\text{COO}^-]$, $[\text{H}^+]$, $[\text{ClCH}_2\text{COOH}]$, and K_a for chloroacetic acid.
- 16.56** A 0.100 M solution of bromoacetic acid (BrCH_2COOH) is 13.2% ionized. Calculate $[\text{H}^+]$, $[\text{BrCH}_2\text{COO}^-]$, and $[\text{BrCH}_2\text{COOH}]$.
- 16.57** A particular sample of vinegar has a pH of 2.90. If acetic acid is the only acid that vinegar contains ($K_a = 1.8 \times 10^{-5}$), calculate the concentration of acetic acid in the vinegar.
- 16.58** How many moles of HF ($K_a = 6.8 \times 10^{-4}$) must be present in 0.200 L to form a solution with a pH of 3.25?
- 16.59** The acid-dissociation constant for benzoic acid ($\text{C}_6\text{H}_5\text{COOH}$) is 6.3×10^{-5} . Calculate the equilibrium concentrations of H_3O^+ , $\text{C}_6\text{H}_5\text{COO}^-$, and $\text{C}_6\text{H}_5\text{COOH}$ in the solution if the initial concentration of $\text{C}_6\text{H}_5\text{COOH}$ is 0.050 M.
- 16.60** The acid-dissociation constant for hypochlorous acid (HClO) is 3.0×10^{-8} . Calculate the concentrations of H_3O^+ , ClO^- , and HClO at equilibrium if the initial concentration of HClO is 0.0090 M.
- 16.61** Calculate the pH of each of the following solutions (K_a and K_b values are given in Appendix D): (a) 0.095 M propionic acid ($\text{C}_2\text{H}_5\text{COOH}$), (b) 0.100 M hydrogen chromate ion (HCrO_4^-), (c) 0.120 M pyridine ($\text{C}_5\text{H}_5\text{N}$).
- 16.62** Determine the pH of each of the following solutions (K_a and K_b values are given in Appendix D): (a) 0.095 M hypochlorous acid, (b) 0.0085 M phenol, (c) 0.095 M hydroxylamine.
- 16.63** Saccharin, a sugar substitute, is a weak acid with $\text{p}K_a = 2.32$ at 25 °C. It ionizes in aqueous solution as follows:
- $$\text{HNC}_7\text{H}_4\text{SO}_3(\text{aq}) \rightleftharpoons \text{H}^+(\text{aq}) + \text{NC}_7\text{H}_4\text{SO}_3^-(\text{aq})$$
- What is the pH of a 0.10 M solution of this substance?
- 16.64** The active ingredient in aspirin is acetylsalicylic acid ($\text{HC}_9\text{H}_7\text{O}_4$), a monoprotic acid with $K_a = 3.3 \times 10^{-4}$ at 25 °C. What is the pH of a solution obtained by dissolving two extra-strength aspirin tablets, containing 500 mg of acetylsalicylic acid each, in 250 mL of water?
- 16.65** Calculate the percent ionization of hydrazoic acid (HN_3) in solutions of each of the following concentrations (K_a is given in Appendix D): (a) 0.400 M, (b) 0.100 M, (c) 0.0400 M.
- 16.66** Calculate the percent ionization of propionic acid ($\text{C}_2\text{H}_5\text{COOH}$) in solutions of each of the following concentrations (K_a is given in Appendix D): (a) 0.250 M, (b) 0.0800 M, (c) 0.0200 M.
- 16.67** Show that for a weak acid, the percent ionization should vary as the inverse square root of the acid concentration.
- 16.68** For solutions of a weak acid, a graph of pH versus the log of the initial acid concentration should be a straight line. What is the magnitude of the slope of that line?
- 16.69** Citric acid, which is present in citrus fruits, is a triprotic acid (Table 16.3). Calculate the pH and the citrate ion ($\text{C}_6\text{H}_5\text{O}_7^{3-}$) concentration for a 0.050 M solution of citric acid. Explain any approximations or assumptions that you make in your calculations.
- 16.70** Tartaric acid is found in many fruits, including grapes, and is partially responsible for the dry texture of certain wines. Calculate the pH and the tartarate ion ($\text{C}_4\text{H}_4\text{O}_6^{2-}$) concentration for a 0.250 M solution of tartaric acid, for which the acid-dissociation constants are listed in Table 16.3. Explain any approximations or assumptions that you make in your calculation.

Weak Bases

- 16.71** What is the essential structural feature of all Brønsted-Lowry bases?
- 16.72** What are two kinds of molecules or ions that commonly function as weak bases?
- 16.73** Write the chemical equation and the K_b expression for the ionization of each of the following bases in aqueous solution: (a) dimethylamine, $(\text{CH}_3)_2\text{NH}$; (b) carbonate ion, CO_3^{2-} ; (c) formate ion, CHO_2^- .
- 16.74** Write the chemical equation and the K_b expression for the reaction of each of the following bases with water: (a) propylamine, $\text{C}_3\text{H}_7\text{NH}_2$; (b) monohydrogen phosphate ion, HPO_4^{2-} ; (c) benzoate ion, $\text{C}_6\text{H}_5\text{CO}_2^-$.
- 16.75** Calculate the molar concentration of OH^- ions in a 0.075 M solution of ethylamine ($\text{C}_2\text{H}_5\text{NH}_2$; $K_b = 6.4 \times 10^{-4}$). Calculate the pH of this solution.
- 16.76** Calculate the molar concentration of OH^- ions in a 0.550 M solution of hypobromite ion (BrO^- ; $K_b = 4.0 \times 10^{-6}$). What is the pH of this solution?
- 16.77** Ephedrine, a central nervous system stimulant, is used in nasal sprays as a decongestant. This compound is a weak organic base:
- $$\text{C}_{10}\text{H}_{15}\text{ON}(\text{aq}) + \text{H}_2\text{O}(\text{l}) \rightleftharpoons \text{C}_{10}\text{H}_{15}\text{ONH}^+(\text{aq}) + \text{OH}^-(\text{aq})$$
- A 0.035 M solution of ephedrine has a pH of 11.33. (a) What are the equilibrium concentrations of $\text{C}_{10}\text{H}_{15}\text{ON}$, $\text{C}_{10}\text{H}_{15}\text{ONH}^+$, and OH^- ? (b) Calculate K_b for ephedrine.
- 16.78** Codeine ($\text{C}_{18}\text{H}_{21}\text{NO}_3$) is a weak organic base. A 5.0×10^{-3} M solution of codeine has a pH of 9.95. Calculate the value of K_b for this substance. What is the $\text{p}K_b$ for this base?

The K_a – K_b Relationship; Acid–Base Properties of Salts

- 16.79** Although the acid-dissociation constant for phenol (C_6H_5OH) is listed in Appendix D, the base-dissociation constant for the phenolate ion ($C_6H_5O^-$) is not. (a) Explain why it is not necessary to list both K_a for phenol and K_b for the phenolate ion. (b) Calculate K_b for the phenolate ion. (c) Is the phenolate ion a weaker or stronger base than ammonia?
- 16.80** We can calculate K_b for the carbonate ion if we know the K_a values of carbonic acid (H_2CO_3). (a) Is K_{a1} or K_{a2} of carbonic acid used to calculate K_b for the carbonate ion? Explain. (b) Calculate K_b for the carbonate ion. (c) Is the carbonate ion a weaker or stronger base than ammonia?
- 16.81** (a) Given that K_a for acetic acid is 1.8×10^{-5} and that for hypochlorous acid is 3.0×10^{-8} , which is the stronger acid? (b) Which is the stronger base, the acetate ion or the hypochlorite ion? (c) Calculate K_b values for CH_3COO^- and ClO^- .
- 16.82** (a) Given that K_b for ammonia is 1.8×10^{-5} and that for hydroxylamine is 1.1×10^{-8} , which is the stronger base? (b) Which is the stronger acid, the ammonium ion or the hydroxylammonium ion? (c) Calculate K_a values for NH_4^+ and H_3NOH^+ .
- 16.83** Using data from Appendix D, calculate $[OH^-]$ and pH for each of the following solutions: (a) 0.10 M NaCN, (b) 0.080 M Na_2CO_3 , (c) a mixture that is 0.10 M in $NaNO_2$ and 0.20 M in $Ca(NO_2)_2$.
- 16.84** Using data from Appendix D, calculate $[OH^-]$ and pH for each of the following solutions: (a) 0.105 M NaF, (b) 0.035 M Na_2S , (c) a mixture that is 0.045 M in CH_3COONa and 0.055 M in $(CH_3COO)_2Ba$.
- 16.85** Predict whether aqueous solutions of the following compounds are acidic, basic, or neutral: (a) NH_4Br , (b) $FeCl_3$, (c) Na_2CO_3 , (d) $KClO_4$, (e) $NaHC_2O_4$.
- 16.86** Predict whether aqueous solutions of the following substances are acidic, basic, or neutral: (a) $CrBr_3$, (b) LiI , (c) K_3PO_4 , (d) $[CH_3NH_3]Cl$, (e) $KHSO_4$.
- 16.87** An unknown salt is either NaF, NaCl, or NaOCl. When 0.050 mol of the salt is dissolved in water to form 0.500 L of solution, the pH of the solution is 8.08. What is the identity of the salt?
- 16.88** An unknown salt is either KBr, NH_4Cl , KCN, or K_2CO_3 . If a 0.100 M solution of the salt is neutral, what is the identity of the salt?
- 16.89** Sorbic acid (C_5H_7COOH) is a weak monoprotic acid with $K_a = 1.7 \times 10^{-5}$. Its salt (potassium sorbate) is added to cheese to inhibit the formation of mold. What is the pH of a solution containing 11.25 g of potassium sorbate in 1.75 L of solution?
- 16.90** Trisodium phosphate (Na_3PO_4) is available in hardware stores as TSP and is used as a cleaning agent. The label on a box of TSP warns that the substance is very basic (caustic or alkaline). What is the pH of a solution containing 35.0 g of TSP in a liter of solution?

Acid–Base Character and Chemical Structure

- 16.91** How does the acid strength of an oxyacid depend on (a) the electronegativity of the central atom; (b) the number of nonprotonated oxygen atoms in the molecule?
- 16.92** (a) How does the strength of an acid vary with the polarity and strength of the $H-X$ bond? (b) How does the acidity of the binary acid of an element vary as a function of the electronegativity of the element? How does this relate to the position of the element in the periodic table?
- 16.93** Explain the following observations: (a) HNO_3 is a stronger acid than HNO_2 ; (b) H_2S is a stronger acid than H_2O ; (c) H_2SO_4 is a stronger acid than HSO_4^- ; (d) H_2SO_4 is a stronger acid than H_2SeO_4 ; (e) CCl_3COOH is a stronger acid than CH_3COOH .
- 16.94** Explain the following observations: (a) HCl is a stronger acid than H_2S ; (b) H_3PO_4 is a stronger acid than H_3AsO_4 ; (c) $HBrO_3$ is a stronger acid than $HBrO_2$; (d) $H_2C_2O_4$ is a stronger acid than $HC_2O_4^-$; (e) benzoic acid (C_6H_5COOH) is a stronger acid than phenol (C_6H_5OH).
- 16.95** Based on their compositions and structures and on conjugate acid–base relationships, select the stronger base in each of the following pairs: (a) BrO^- or ClO^- , (b) BrO^- or BrO_2^- , (c) HPO_4^{2-} or $H_2PO_4^-$.
- 16.96** Based on their compositions and structures and on conjugate acid–base relationships, select the stronger base in each of the following pairs: (a) NO_3^- or NO_2^- , (b) PO_4^{3-} or AsO_4^{3-} , (c) HCO_3^- or CO_3^{2-} .
- 16.97** Indicate whether each of the following statements is true or false. For each statement that is false, correct the statement to make it true. (a) In general, the acidity of binary acids increases from left to right in a given row of the periodic table. (b) In a series of acids that have the same central atom, acid strength increases with the number of hydrogen atoms bonded to the central atom. (c) Hydrotelluric acid (H_2Te) is a stronger acid than H_2S because Te is more electronegative than S.

16.98 Indicate whether each of the following statements is true or false. For each statement that is false, correct the statement to make it true. (a) Acid strength in a series of H—X molecules increases with increasing size of X. (b) For acids of the same general structure


but differing electronegativities of the central atoms, acid strength decreases with increasing electronegativity of the central atom. (c) The strongest acid known is HF because fluorine is the most electronegative element.

Lewis Acids and Bases


16.99 If a substance is an Arrhenius base, is it necessarily a Brønsted–Lowry base? Is it necessarily a Lewis base? Explain.

16.100 If a substance is a Lewis acid, is it necessarily a Brønsted–Lowry acid? Is it necessarily an Arrhenius acid? Explain.

16.101 Identify the Lewis acid and Lewis base among the reactants in each of the following reactions:



16.102 Identify the Lewis acid and Lewis base in each of the following reactions:



16.103 Predict which member of each pair produces the more acidic aqueous solution: (a) K^{+} or Cu^{2+} , (b) Fe^{2+} or Fe^{3+} , (c) Al^{3+} or Ga^{3+} . Explain.

16.104 Which member of each pair produces the more acidic aqueous solution: (a) ZnBr_2 or CdCl_2 , (b) CuCl or $\text{Cu}(\text{NO}_3)_2$, (c) $\text{Ca}(\text{NO}_3)_2$ or NiBr_2 ? Explain.


ADDITIONAL EXERCISES

16.105 In your own words, define or explain (a) K_w , (b) K_a , (c) pOH , (d) $\text{p}K_b$.

16.106 Indicate whether each of the following statements is correct or incorrect. For those that are incorrect, explain why they are wrong.

- (a) Every Brønsted–Lowry acid is also a Lewis acid.
 (b) Every Lewis acid is also a Brønsted–Lowry acid.
 (c) Conjugate acids of weak bases produce more acidic solutions than conjugate acids of strong bases.
 (d) K^{+} ion is acidic in water because it causes hydrating water molecules to become more acidic.
 (e) The percent ionization of a weak acid in water increases as the concentration of acid decreases.


16.107 Predict whether the equilibrium lies to the right or to the left in the following reactions:



16.108 The odor of fish is due primarily to amines, especially methylamine (CH_3NH_2). Fish is often served with a wedge of lemon, which contains citric acid. The amine and the acid react forming a product with no odor, thereby making the less-than-fresh fish more appetizing.

Using data from Appendix D, calculate the equilibrium constant for the reaction of citric acid with methylamine, if only the first proton of the citric acid (K_{a1}) is important in the neutralization reaction.

16.109 Hemoglobin plays a part in a series of equilibria involving protonation-deprotonation and oxygenation-deoxygenation. The overall reaction is approximately as follows:



where Hb stands for hemoglobin, and HbO_2 for oxyhemoglobin. (a) The concentration of O_2 is higher in the lungs and lower in the tissues. What effect does high $[\text{O}_2]$ have on the position of this equilibrium? (b) The normal pH of blood is 7.4. Is the blood acidic, basic, or neutral? (c) If the blood pH is lowered by the presence of large amounts of acidic metabolism products, a condition known as acidosis results. What effect does lowering blood pH have on the ability of hemoglobin to transport O_2 ?

16.110 Calculate the pH of a solution made by adding 2.50 g of lithium oxide (Li_2O) to enough water to make 1.500 L of solution.

16.111 Which of the following solutions has the higher pH? (a) a 0.1 M solution of a strong acid or a 0.1 M solution of a weak acid, (b) a 0.1 M solution of an acid with $K_a = 2 \times 10^{-3}$ or one with $K_a = 8 \times 10^{-6}$, (c) a 0.1 M solution of a base with $\text{p}K_b = 4.5$ or one with $\text{p}K_b = 6.5$.

- [16.112] What is the pH of a solution that is $2.5 \times 10^{-9} M$ in NaOH? Does your answer make sense?
- 16.113** Caproic acid ($C_5H_{11}COOH$) is found in small amounts in coconut and palm oils and is used in making artificial flavors. A saturated solution of the acid contains 11 g/L and has a pH of 2.94. Calculate K_a for the acid.
- [16.114] A hypothetical acid H_2X is both a strong acid and a diprotic acid. (a) Calculate the pH of a 0.050 M solution of H_2X , assuming that only one proton ionizes per acid molecule. (b) Calculate the pH of the solution from part (a), now assuming that both protons of each acid molecule completely ionize. (c) In an experiment it is observed that the pH of a 0.050 M solution of H_2X is 1.27. Comment on the relative acid strengths of H_2X and HX^- . (d) Would a solution of the salt $NaHX$ be acidic, basic, or neutral? Explain.
- 16.115** Butyric acid is responsible for the foul smell of rancid butter. The pK_a of butyric acid is 4.84. (a) Calculate the pK_b for the butyrate ion. (b) Calculate the pH of a 0.050 M solution of butyric acid. (c) Calculate the pH of a 0.050 M solution of sodium butyrate.
- 16.116** Arrange the following 0.10 M solutions in order of increasing acidity (decreasing pH): (i) NH_4NO_3 , (ii) $NaNO_3$, (iii) CH_3COONH_4 , (iv) NaF , (v) CH_3COONa .
- [16.117] What are the concentrations of H^+ , $H_2PO_4^-$, HPO_4^{2-} , and PO_4^{3-} in a 0.0250 M solution of H_3PO_4 ?
- [16.118] Many moderately large organic molecules containing basic nitrogen atoms are not very soluble in water as neutral molecules, but they are frequently much more soluble as their acid salts. Assuming that pH in the stomach is 2.5, indicate whether each of the following compounds would be present in the stomach as the neutral base or in the protonated form: nicotine, $K_b = 7 \times 10^{-7}$; caffeine, $K_b = 4 \times 10^{-14}$; strychnine, $K_b = 1 \times 10^{-6}$; quinine, $K_b = 1.1 \times 10^{-6}$.
- [16.119] The amino acid glycine (H_2N-CH_2-COOH) can participate in the following equilibria in water:
- $$H_2N-CH_2-COOH + H_2O \rightleftharpoons H_2N-CH_2-COO^- + H_3O^+ \quad K_a = 4.3 \times 10^{-3}$$
- $$H_2N-CH_2-COOH + H_2O \rightleftharpoons {}^+H_3N-CH_2-COOH + OH^- \quad K_b = 6.0 \times 10^{-5}$$
- (a) Use the values of K_a and K_b to estimate the equilibrium constant for the intramolecular proton transfer to form a zwitterion:
- $$H_2N-CH_2-COOH \rightleftharpoons {}^+H_3N-CH_2-COO^-$$
- What assumptions did you need to make? (b) What is the pH of a 0.050 M aqueous solution of glycine? (c) What would be the predominant form of glycine in a solution with pH 13? With pH 1?
- 16.120** The structural formula for acetic acid is shown in Table 16.2. Replacing hydrogen atoms on the carbon with chlorine atoms causes an increase in acidity, as follows:


Acid	Formula	K_a (25 °C)
Acetic	CH_3COOH	1.8×10^{-5}
Chloroacetic	$CH_2ClCOOH$	1.4×10^{-3}
Dichloroacetic	$CHCl_2COOH$	3.3×10^{-2}
Trichloroacetic	CCl_3COOH	2×10^{-1}

Using Lewis structures as the basis of your discussion, explain the observed trend in acidities in the series. Calculate the pH of a 0.010 M solution of each acid.

INTEGRATIVE EXERCISES



- 16.121** Calculate the number of $H^+(aq)$ ions in 1.0 mL of pure water at 25 °C.
- 16.122** How many milliliters of concentrated hydrochloric acid solution (36.0% HCl by mass, density = 1.18 g/mL) are required to produce 10.0 L of a solution that has a pH of 2.05?
- 16.123** The volume of an adult's stomach ranges from about 50 mL when empty to 1 L when full. If the stomach volume is 400 mL and its contents have a pH of 2, how many moles of H^+ does the stomach contain? Assuming that all the H^+ comes from HCl, how many grams of sodium hydrogen carbonate will totally neutralize the stomach acid?
- 16.124** Atmospheric CO_2 levels have risen by nearly 20% over the past 40 years from 315 ppm to 380 ppm. (a) Given that the average pH of clean, unpolluted rain today is 5.4, determine the pH of unpolluted rain 40 years ago. Assume that carbonic acid (H_2CO_3) formed by the reaction of CO_2 and water is the only factor influencing pH.
- $$CO_2(g) + H_2O(l) \rightleftharpoons H_2CO_3(aq)$$
- (b) What volume of CO_2 at 25 °C and 1.0 atm is dissolved in a 20.0-L bucket of today's rainwater?
- [16.125] In many reactions the addition of $AlCl_3$ produces the same effect as the addition of H^+ . (a) Draw a Lewis structure for $AlCl_3$ in which no atoms carry formal charges, and determine its structure using the VSEPR method. (b) What characteristic is notable about the structure in part (a) that helps us understand the acidic character of $AlCl_3$? (c) Predict the result of the reaction between $AlCl_3$ and NH_3 in a solvent that does not participate as a reactant. (d) Which acid-base theory is most suitable for discussing the similarities between $AlCl_3$ and H^+ ?
- [16.126] What is the boiling point of a 0.10 M solution of $NaHSO_4$ if the solution has a density of 1.002 g/mL?
- [16.127] Cocaine is a weak organic base whose molecular formula is $C_{17}H_{21}NO_4$. An aqueous solution of cocaine was found to have a pH of 8.53 and an osmotic pressure of 52.7 torr at 15 °C. Calculate K_b for cocaine.

[16.128] The iodate ion is reduced by sulfite according to the following reaction:



The rate of this reaction is found to be first order in IO_3^- , first order in SO_3^{2-} , and first order in H^+ . (a) Write the rate law for the reaction. (b) By what factor will the rate of the reaction change if the pH is lowered from 5.00 to 3.50? Does the reaction proceed faster or slower at the lower pH? (c) By using the concepts discussed in Section 14.6, explain how the reaction can be pH-dependent even though H^+ does not appear in the overall reaction.

[16.129] (a) Using dissociation constants from Appendix D, determine the value for the equilibrium constant for each of the following reactions. (Remember that when reactions are added, the corresponding equilibrium constants are multiplied.)



(b) We usually use single arrows for reactions when the forward reaction is appreciable (K much greater than 1) or when products escape from the system, so that equilibrium is never established. If we follow this convention, which of these equilibria might be written with a single arrow?

[16.130] Lactic acid, $\text{CH}_3\text{CH}(\text{OH})\text{COOH}$, received its name because it is present in sour milk as a product of bacterial action. It is also responsible for the soreness in muscles after vigorous exercise. (a) The $\text{p}K_a$ of lactic acid is 3.85. Compare this with the value for propionic acid ($\text{CH}_3\text{CH}_2\text{COOH}$, $\text{p}K_a = 4.89$), and explain the difference. (b) Calculate the lactate ion concentration in a 0.050 M solution of lactic acid. (c) When a solution of sodium lactate, $\text{CH}_3\text{CH}(\text{OH})\text{COONa}$, is mixed with an aqueous copper(II) solution, it is possible to obtain a solid salt of copper(II) lactate as a blue-green hydrate, $(\text{CH}_3\text{CH}(\text{OH})\text{COO})_2\text{Cu} \cdot x\text{H}_2\text{O}$. Elemental analysis of the solid tells us that the solid is 22.9% Cu and 26.0% C by mass. What is the value for x in the formula for the hydrate? (d) The acid-dissociation constant for the $\text{Cu}^{2+}(aq)$ ion is 1.0×10^{-8} . Based on this value and the acid-dissociation constant of lactic acid, predict whether a solution of copper(II) lactate will be acidic, basic, or neutral. Explain your answer.